### For Research Use Only

# NDP52 Polyclonal antibody

Catalog Number:12229-1-AP

Featured Product

**44 Publications** 



**Basic Information** 

Catalog Number:

12229-1-AP

Size:

GeneID (NCBI):

600 µg/ml

10241

Source:

Rabbit

Q13137

Isotype:

GeneID (NCBI):

GeneID (NCBI):

Full Name:

gG calcium binding and coiled-coil

Immunogen Catalog Number: domain 2

AG2866 Calculated MW:

446 aa, 52 kDa Observed MW: 52 kDa

Observed MW

**Applications** 

Tested Applications: IF/ICC, IHC, IP, WB, ELISA Cited Applications:

IF, IP, WB

Species Specificity: human, mouse, rat Cited Species:

human, rat, mouse, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method: Antigen affinity purification

Recommended Dilutions: WB 1:500-1:2000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500 IF 1:200-1:800

**Positive Controls:** 

WB: HeLa cells, rat skeletal muscle tissue

P: HeLa cells,

IHC: human ovary tumor tissue, human skeletal

muscle tissue

IF : HeLa cells,

# **Background Information**

NDP52, also named as CALCOCO2, is an autophagy receptor. It plays a role in ruffle formation and actin cytoskeleton organization. Mouse/Rat NDP52 has some isoforms with MW 28-40 kDa and 67 kDa. Human NDP52 has some isoforms with MW 43-47 kDa and 52-55 kDa,

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Shouheng Jin	28965816	Mol Cell	WB
Sebastian Boland	36207292	Nat Commun	WB
Jureepon Roboon	36274991	Front Cell Neurosci	WB

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using NDP52 antibody (12229-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




HeLa cells were subjected to SDS PAGE followed by western blot with 12229-1-AP (NDP52 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours




IP result of anti-NDP52 (IP:12229-1-AP, 3ug; Detection:12229-1-AP 1:1000) with HeLa cells lysate 2150ug.



rat skeletal muscle tissue were subjected to SDS PAGE followed by western blot with 12229-1-AP (NDP52 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 12229-1-AP (NDP52 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 12229-1-AP (NDP52 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).