For Research Use Only

ZFP36L1/2 Polyclonal antibody

Catalog Number:12306-1-AP

Featured Product


5 Publications




Basic Information	Catalog Number: 12306-1-AP	5		Purification Method: Antigen affinity purification	
	-			Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500 IF 1:20-1:200	
	400 µg/ml				
	Isotype:Full Name:IgGzinc finger protein 36, C3H type-like 1		xe 1		
	Immunogen Catalog Number: AG2952		Calculated MW: 338 aa, 36 kDa Observed MW: 45-50 kDa		
Applications	Tested Applications: IF/ICC, IHC, WB, ELISA		Positive Controls:		
	Cited Applications:	WD. 0-937		37 cells, 3T3-L1 cells, RAW 264.7 cells, HeLa	
	IF, IHC, WB			ary tissue,	
	Species Specificity: human, mouse, rat		IF : HepG2 cells,		
	Cited Species: human, mouse				
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	ZFP36L1, also named as Butyrate response factor 1 or TPA-induced sequence 11b, is a 338 amino acid protein, which is phosphorylated by RPS6KA1 at Ser-334 upon phorbol 12-myristate 13-acetate (PMA) treatment. ZFP36 encodes tristetraprolin (TTP) the prototype of a small family of RBPs, called the ZFP36 family, that are characterised by highly conserved tandem CCCH zinc-finger RNA-binding domains [PMID: 10751406]. ZFP36 is a RBP that promotes RNA decay and negatively regulates the expression of the myogenic regulatory factor MyoD by binding to the 3'UTR of MyoD and NRNA [PMID: 25815583]. Mouse satellite cells from Zfp36-deficient mice express increased amounts of MyoD and display impaired satellite activation, demonstrating a role for ZFP36 in the maintenance of quiescence [PMID: 25815583]. The functions of the ZFP36L1 and ZFP36L2 family members have not been evaluated in skeletal muscle stem cell fate, but have been shown to act redundantly to promote quiescence during lymphocyte development. ZFP36L1 has been implicated in the persistence of the marginal zone B lymphocyte population. ZFP36L1 exists three isoform through blasting on NCBI database and can be phosphorylated, so the range of the molecular weight of ZFP36L1 is about 40-50 kDa. (PMID: 26180518, PMID: 17030620). This antibody recognizes both ZFP36L1 and ZFP36L2.				
Notable Publications	Author	Pubmed ID	Journal	Application	
	Weirui Ma	30449617	Cell	IF	
	Xiao Feng	33473330	Mol Ther Nucleic A	cids IF	
	Fang Wu	34131106	Cell Death Dis	WB,IHC,IF	
Storage	Storage: Store at -20°C. Stable for one yea Storage Buffer: PBS with 0.02% sodium azide ar Aliquoting is unnecessary for -20	nd 50% glycerol pH	7.3.		

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




U-937 cells were subjected to SDS PAGE followed by western blot with 12306-1-AP (ZFP36L1/2 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.




WB result of ZFP36L1/2 antibody (12306-1-AP; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-ZFP36L1/2 transfected HeLa cells.



Immunohistochemical analysis of paraffinembedded rat ovary tissue slide using 12306-1-AP (ZFP36L1/2 antibody) at dilution of 1:100 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat ovary tissue slide using 12306-1-AP (ZFP36L1/2 antibody) at dilution of 1:100 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of HepG2 cells using 12306-1-AP (ZFP36L1/2 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).