

SMAD2 Polyclonal antibody

Catalog Number: 12570-1-AP

Featured Product

188 Publications

Basic Information

Catalog Number:

12570-1-AP

Concentration:

450 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG3237

GenBank Accession Number:

BC014840

GeneID (NCBI):

4087

UNIPROT ID:

Q15796

Full Name:

SMAD family member 2

Calculated MW:

467 aa, 52 kDa

Observed MW:

52-70 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB: 1:2000-1:10000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC: 1:50-1:500

IF/ICC: 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IHC, IF, IP, CoIP, ChIP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A549 cells, HeLa cells, mouse skeletal muscle tissue, rat skeletal muscle tissue, HEK-293, HepG2 cells, MCF-7 cells, PC-3 cells, C6 cells, HEK-293 cells, HT-1080 cells, HUVEC cells, C2C12 cells

IP: HepG2 cells,

IHC: human colon cancer tissue, human stomach cancer tissue, human endometrial cancer tissue, mouse colon tissue, rat colon tissue

IF/ICC: HepG2 cells,

Background Information

SMAD2, also named as MADH2 and MADR2, belongs to the dwarfin/SMAD family, contains 1 MH1 (MAD homology 1) domain and 1 MH2 (MAD homology 2) domain. SMAD2 is a receptor-regulated SMAD (R-SMAD) that is an intracellular signal transducer and transcriptional modulator activated by TGF-beta (transforming growth factor) and activin type 1 receptor kinases. This protein may act as a tumor suppressor in colorectal carcinoma. It is phosphorylated on one or several of Thr-220, Ser-245, Ser-250, and Ser-255. In response to TGF-beta, it is phosphorylated on Ser-465/467 by TGF-beta and activin type 1 receptor kinases, and then able to interact with SMURF2, recruiting other proteins, such as SNON, for degradation. In response to decorin, the naturally occurring inhibitor of TGF-beta signaling, it is phosphorylated on Ser-240 by CaMK2. It is phosphorylated by MAPK3 upon EGF stimulation; which increases transcriptional activity and stability, and is blocked by calmodulin. In response to TGF-beta, it is ubiquitinated by NEDD4L, which promotes its degradation. In response to TGF-beta signaling, it is acetylated on Lys-19 by coactivators, which increases transcriptional activity. This antibody is a rabbit polyclonal antibody raised against residues near the N terminus of human SMAD2. The molecular weight of unphosphorylated forms of Smad2 is 52 kDa and phosphorylated forms of Smad2 is 58 kDa. (PMID: 9006934). The ubiquitination form of Smad2 is ~70 kDa (PMID: 25998442).

Notable Publications

Author	Pubmed ID	Journal	Application
Shun Gu	33007305	Exp Eye Res	WB
Shaling Li	36169092	Cancer Sci	WB
Bingyu Xie	36179941	Mol Cell Endocrinol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 12570-1-AP (SMAD2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




IP result of anti-SMAD2 (IP:12570-1-AP, 3ug; Detection:12570-1-AP 1:1000) with HepG2 cells lysate 3000ug.



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 12570-1-AP (SMAD2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 12570-1-AP (SMAD2 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using SMAD2 antibody (12570-1-AP) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).