## For Research Use Only

# ILK Polyclonal antibody

Catalog Number: 12955-1-AP

7 Publications



**Basic Information** 

Catalog Number: 12955-1-AP Size:

450 µg/ml Source:

Rabbit Isotype:

Immunogen Catalog Number:

AG3570

Observed MW:

GenBank Accession Number:

BC001554 GeneID (NCBI):

**UNIPROT ID:** Q13418 Full Name:

integrin-linked kinase

Calculated MW: 451 aa, 59 kDa

51-59 kDa

**Purification Method:** 

Antigen affinity purification

Recommended Dilutions: WB 1:500-1:2000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

**Applications** 

**Tested Applications:** IHC, IP, WB,ELISA Cited Applications:

WB, IP, IF, IHC

Species Specificity: human, mouse, rat **Cited Species:** human, mouse, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

**Positive Controls:** 

WB: mouse brain tissue, mouse kidney tissue

IP: mouse kidney tissue,

IHC: human prostate cancer tissue,

# **Background Information**

Integrin-linked kinase (ILK) is an ankyrin-repeat containing serine-threonine protein kinase capable of interacting with the cytoplasmic domains of integrin  $\beta$  1,  $\beta$  2, and  $\beta$  3 subunits and whose kinase activity is modulated by cell $extracellular\ matrix\ interactions\ (PMID: 9539744).\ This\ protein\ belongs\ to\ the\ protein\ kinase\ superfamily.\ It\ can\ be$ autophosphorylated on serine residues.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Lan Lin	32485356	Int Immunopharmacol	WB
Ying Feng	33531984	J Cancer	WB
Q Zhou	30739146	Osteoporos Int	WB

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 12955-1-AP (ILK antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 12955-1-AP (ILK antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-ILK (IP:12955-1-AP, 4ug; Detection:12955-1-AP 1:800) with mouse kidney tissue lysate 4000ug.