### For Research Use Only

# CPT1C-specific Polyclonal antibody

Catalog Number: 12969-1-AP

10 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 12969-1-AP BC029104 GeneID (NCBI): Size: 550 μg/ml 126129 **UNIPROT ID:** Source: Rabbit Q8TCG5 Full Name: Isotype:

carnitine palmitoyltransferase 1C

Calculated MW: Immunogen Catalog Number: AG3649 792 aa, 90 kDa Observed MW:

70-82 kDa

**Purification Method:** Antigen affinity purification Recommended Dilutions:

WB 1:500-1:1000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

**Applications** 

**Tested Applications:** IHC, IP, WB, ELISA **Cited Applications:** WB. IF

Species Specificity: human, mouse, rat **Cited Species:** human, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

WB: mouse brain tissue, HeLa cells, rat brain tissue

IP: mouse testis tissue,

IHC: human gliomas tissue, human testis tissue,

human brain tissue

## **Background Information**

CPT1C belongs to the carnitine/choline acetyltransferase family. Carnitine palmitoyltransferase (CPT) deficiencies are common disorders of mitochondrial fatty acid oxidation. The CPT system is made up of two separate proteins located in the outer (CPT1) and inner (CPT2) mitochondrial membranes. CPT1C is an active forms of related braintype carnitine palmitoyltransferase I. CPT1C may be a regulated target of malonyl-CoA that relays the "malonyl-CoA signal" in hypothalamic neurons that express the orexigenic and anorexigenic neuropeptides that regulate food intake and peripheral energy expenditure (PMID: 17018521). This antibody specifically recognizes CPT1C, and does not cross-react with CPT1A or CPT1B.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
De Huang	25242319	Cell Rep	WB
Carlo Rinaldi	25751282	JAMA Neurol	IF
Brenda Raud	30043753	Cell Metab	WB

Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## **Selected Validation Data**




mouse brain tissue were subjected to SDS PAGE followed by western blot with 12969-1-AP (CPT1C-specific antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



IP result of anti-CPT1C-specific (IP:12969-1-AP, 5ug; Detection:12969-1-AP 1:500) with mouse testis tissue lysate 8000ug.



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 12969-1-AP (CPT1C-specific antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 12969-1-AP (CPT1C-specific antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).