For Research Use Only

CNPase Polyclonal antibody

Catalog Number: 13427-1-AP

15 Publications



Basic Information

Catalog Number: 13427-1-AP Size: 550 μg/ml Source: Rabbit

Immunogen Catalog Number:

AG4227

Isotype:

GenBank Accession Number:

BC028040 GeneID (NCBI): 1267 **UNIPROT ID:** P09543

2',3'-cyclic nucleotide 3' phosphodiesterase Calculated MW:

421 aa, 48 kDa Observed MW: 45-48 kDa

Full Name:

Purification Method: Antigen affinity purification

Recommended Dilutions: WB 1:500-1:1000 IHC 1:20-1:200

IF 1:50-1:500

Applications

Tested Applications: FC, IF/ICC, IHC, WB, ELISA

IF, IHC, WB Species Specificity: human, mouse, rat **Cited Species:**

human, rat, mouse

Cited Applications:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HepG2 cells,

IHC: mouse brain tissue, IF: mouse brain tissue,

Background Information

CNP(2',3'-cyclic-nucleotide 3'-phosphodiesterase) belongs to the cyclic nucleotide phosphodiesterase family. It plays an essential role in axonal survival but not in myelin assembly, coupling oligodendroglial functions in axonal support and myelination. This protein has 2 isoforms produced by alternative splicing with molecular mass of 45

Notable Publications

Author	Pubmed ID	Journal	Application
Jiuyang Ding	34562559	Toxicol Lett	IHC
Li Zhang	32921051	Acta Physiol (Oxf)	WB
Ting Gao	36460235	Neurosci Lett	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using 13427-1-AP (CNPase antibody) at dilution of 1:50 and Alexa Fluor 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using 13427-1-AP (CNPase antibody) at dilution of 1:50 and Alexa Fluor 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




HepG2 cells were subjected to SDS PAGE followed by western blot with 13427-1-AP (CNPase antibody at dilution of 1:600 incubated at room temperature for 1.5 hours.



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug Anti-Human CNPase (13427-1-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 13427-1-AP (CNPase antibody at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 13427-1-AP (CNPase antibody at dilution of 1:50 (under 40x lens).