

Carboxypeptidase E Polyclonal antibody

Catalog Number: 13710-1-AP

Featured Product

8 Publications

Basic Information

Catalog Number:

13710-1-AP

Size:

153 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG4647

GenBank Accession Number:

BC033866

GeneID (NCBI):

1363

UNIPROT ID:

P16870

Full Name:

carboxypeptidase E

Calculated MW:

476 aa, 53 kDa

Observed MW:

50 kDa, 52 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:20-1:200

IF 1:50-1:500

Applications

Tested Applications:

IF/ICC, IHC, IP, WB, ELISA

Cited Applications:

WB, IF, IHC

Species Specificity:

human, mouse, rat

Cited Species:

human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : mouse brain tissue,

IP : mouse brain tissue,

IHC : human pituitary tissue,

IF : HepG2 cells, mouse brain tissue

Background Information

CPE(Carboxypeptidase E) is also named as CPH. It belongs to the peptidase M14 family and is involved in the biosynthesis of peptide hormones and neurotransmitters, including insulin. The deduced 476-amino acid human protein contains an N-terminal signal peptide, followed by a 'pro' sequence, a polyarginine stretch, and 2 potential N-glycosylation sites.

Notable Publications

Author	Pubmed ID	Journal	Application
Kristen E Rohli	36325514	Function (Oxf)	WB
Jing Yu	34630524	Front Genet	WB, IHC
Seokwon Jo	33115825	Diabetes	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human pituitary using 13710-1-AP (Carboxypeptidase E antibody) at dilution of 1:50 (under 40x lens).




Immunohistochemical analysis of paraffin-embedded human pituitary using 13710-1-AP (Carboxypeptidase E antibody) at dilution of 1:50 (under 10x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using Carboxypeptidase E antibody (13710-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).



mouse brain tissue were subjected to SDS PAGE followed by western blot with 13710-1-AP (Carboxypeptidase E antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



IP result of anti-Carboxypeptidase E (IP:13710-1-AP, 3ug; Detection:13710-1-AP 1:500) with mouse brain tissue lysate 3500ug.