## For Research Use Only

## Carboxypeptidase E Polyclonal antibody



Catalog Number: 13710-1-AP

**Featured Product** 

8 Publications

BC033866

1363

P16870

GeneID (NCBI):

**UNIPROT ID:** 

Full Name:

carboxypeptidase E Calculated MW:

GenBank Accession Number:

**Basic Information** 

Catalog Number: 13710-1-AP Size: 153 μg/ml Source:

Rabbit Isotype:

Immunogen Catalog Number:

AG4647

Observed MW:

50 kDa, 52 kDa

476 aa, 53 kDa

**Purification Method:** 

Antigen affinity purification

Recommended Dilutions: WB 1:500-1:1000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:20-1:200 IF 1:50-1:500

**Applications** 

**Tested Applications:** IF/ICC, IHC, IP, WB, ELISA

Cited Applications: IF, IHC, WB Species Specificity:

human, mouse, rat **Cited Species:** human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse brain tissue, IP: mouse brain tissue, IHC: human pituitary tissue, IF: HepG2 cells, mouse brain tissue

**Background Information** 

CPE(Carboxypeptidase E) is also named as CPH.It belongs to the peptidase M14 family and is involved in the biosynthesis of peptide hormones and neurotransmitters, including insulin. The deduced 476-amino acid human protein contains an N-terminal signal peptide, followed by a 'pro' sequence, a polyarginine stretch, and 2 potential N-glycosylation sites.

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Kristen E Rohli	36325514	Function (Oxf)	WB
Jing Yu	34630524	Front Genet	WB,IHC
Seokwon Jo	33115825	Diabetes	WB


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Immunohistochemical analysis of paraffinembedded human pituitary using 13710-1-AP (Carboxypeptidase E antibody) at dilution of 1:50 (under 10x lens).




Immunohistochemical analysis of paraffinembedded human pituitary using 13710-1-AP (Carboxypeptidase E antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using Carboxypeptidase E antibody (13710-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).



mouse brain tissue were subjected to SDS PAGE followed by western blot with 13710-1-AP (Carboxypeptidase E antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



IP result of anti-Carboxypeptidase E (IP:13710-1-AP, 3ug; Detection:13710-1-AP 1:500) with mouse brain tissue lysate 3500ug.