## For Research Use Only

# Calponin Polyclonal antibody

Catalog Number: 13938-1-AP

Featured Product

54 Publications



**Basic Information** 

Catalog Number:

13938-1-AP

BC022015

Concentration:

500 ug/ml

Source:

Rabbit

Cotype:

GenBank Accession Number:

BC022015

GeneID (NCBI):

1264

UNIPROT ID:

P51911

Isotype:

Full Name:

calponin 1, basic, smooth muscle

Immunogen Catalog Number: Calculated MW:

AG4989 33 kDa

Observed MW: 35 kDa Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:5000-1:50000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

protein lysate IHC 1:2000-1:8000 IF-P 1:50-1:500 IF/ICC 1:200-1:800

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, IF-P, FC (Intra), IP, ELISA

Cited Applications: WB, IHC, IF Species Specificity:

human, mouse, rat Cited Species:

human, mouse, rat, pig, bovine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse colon tissue, mouse heart tissue, mouse lung tissue, mouse uterus tissue

IP: mouse lung tissue,

IHC: human appendicitis tissue, human ovary tumor tissue, human hysteromyoma tissue, human breast cancer tissue, human bladder tissue, human skin tissue. human heart tissue

IF-P: mouse heart tissue, IF/ICC: C2C12 cells, FC (Intra): C2C12 cells,

## **Background Information**

Calponin is a family of actin filament-associated proteins which regulate smooth muscle cell contraction. Three isoforms of calponin exist: calponin h1 (CNN1), calponin h2 (CNN2) and calponin 3 (CNN3). CNN1 is a basic 34-kD protein specifically expressed in smooth muscle and a marker of smooth muscle cell differentiation. This antibody raised against the full-length of human CNN1 may cross-react with CNN2 and CNN3.

### **Notable Publications**

Author	Pubmed ID	Journal	Application
Yang Yang	31523025	Acta Biomater	IF
Yan Yan	36068532	Reprod Biol Endocrinol	IHC
Meng Wang	30651790	Exp Ther Med	IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


#### Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 13938-1-AP (Calponin antibody) at dilution of 1:30000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human heart using 13938-1-AP (CNN1 antibody) at dilution of 1:50 (under 40x lens).




Immunohistochemical analysis of paraffinembedded human heart using 13938-1-AP (CNN1 antibody) at dilution of 1:50 (under 10x lens).




mouse heart tissue were subjected to SDS PAGE followed by western blot with 13938-1-AP (CNN1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse heart tissue using Calponin antibody (13938-1-AP) at dilution of 1:200 and Multi-rAb Coralite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (-20°C Methanol) fixed C2C12 cells using Calponin antibody (13938-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




IP result of anti-Calponin (IP:13938-1-AP, 4ug: Detection:13938-1-AP 1:1000) with mouse lung tissue lysate 2800ug.




Immunohistochemical analysis of paraffinembedded human appendicitis tissue slide using 13938-1-AP (Calponin antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse heart tissue using 13938-1-AP (Calponin antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).





Immunofluorescent analysis of (-20°C Methanol) fixed C2C12 cells using Calponin antibody (13938-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).

1X10^6 C2C12 cells were intracellularly stained with 0.4 ug Anti-Human Calponin (13938-1-AP) and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit I gG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).