

For Research Use Only

CD43 Polyclonal antibody

Catalog Number: 13959-1-AP



Basic Information

Catalog Number: 13959-1-AP	GenBank Accession Number: BC012350	Purification Method: Antigen affinity purification
Size: 1000 µg/ml	GeneID (NCBI): 6693	Recommended Dilutions: WB 1:500-1:2000 IHC 1:20-1:200
Source: Rabbit	ENSEMBL Gene ID: ENSG00000197471	
Isotype: IgG	UNIPROT ID: P16150	
Immunogen Catalog Number: AG4836	Full Name: sialophorin	
	Calculated MW: 400 aa, 43 kDa	
	Observed MW: 60-66 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls: WB : K-562 cells, Jurkat cells
Species Specificity: human	IHC : human lymphoma tissue,
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

CD43, also named leukosialin, sialophorin, galactoglycoprotein, leukocyte sialoglycoprotein, SPN, and is a mucin-like type I transmembrane protein. In humans, it is expressed in haematopoietic cells, including T lymphocytes, monocytes, granulocytes, natural killer cells, platelets, and haematopoietic stem cells, with the exception of mature erythrocytes and B cell subpopulations. The unglycosylated form of CD43 migrates on SDS-PAGE with a molecular weight of 54 kDa (PMID: 2943740), whereas CD43 from different haematopoietic cell lines displays increased molecular weights since it is glycosylated with O-linked chains that differ in core structure and sialylation.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



K-562 cells were subjected to SDS PAGE followed by western blot with 13959-1-AP (CD43 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human lymphoma using 13959-1-AP (CD43 antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human lymphoma using 13959-1-AP (CD43 antibody) at dilution of 1:50 (under 40x lens).