For Research Use Only

CNOT7 Polyclonal antibody

Catalog Number:14102-1-AP

Featured Product

6 Publications



Basic Information

Catalog Number: 14102-1-AP

Size: 253 μg/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG5235

Tested Applications:

Cited Applications:

Species Specificity: human, mouse, rat Cited Species:

GenBank Accession Number:

BC060852 GeneID (NCBI): 29883 **UNIPROT ID:** Q9UIV1 Full Name:

CCR4-NOT transcription complex, subunit 7

Calculated MW: 285 aa, 33 kDa Observed MW:

33-35 kDa

Purification Method:

Antigen affinity purification Recommended Dilutions:

WB 1:500-1:1000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate

Applications

IP, WB,ELISA

human, mouse

Positive Controls:

WB: mouse testis tissue, IP: mouse testis tissue,

Background Information

Cnot7, Ccr4-Not complex 7, is a component of CCR4-NOT complex and known to be a transcription factor and a modulator of mRNA degradation in yeast. Mammalian Cnot7 interacts with Tob and its family members, which is known as an inhibitor of BMP signaling through interfering with the Smad system and is involved in bone metabolism [PMID:17451368]. It is one of at least 8 subunits that make up the CCR4-Not transcription complex, which is involved in negative regulation of the pol II directed transcription in yeast 24 and may be involved inglobal gene regulation by modulation of TATAA-binding protein (TBP) activity [PMID:10880468].

Notable Publications

Author	Pubmed ID	Journal	Application
Masashi Yamaji	28297718	Nature	WB
Hsin-Hui Hsieh	34090459	J Inflamm (Lond)	WB
Roy Matkovic	35013187	Nat Commun	IP


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



mouse testis tissue were subjected to SDS PAGE followed by western blot with 14102-1-AP (CNOT7 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



IP result of anti-CNOT7 (IP:14102-1-AP, 4ug; Detection:14102-1-AP 1:300) with mouse testis tissue lysate 4800ug.