## For Research Use Only

# ACAT2 Polyclonal antibody

Catalog Number: 14755-1-AP

**Featured Product** 

8 Publications



Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

WB 1:2000-1:12000

protein lysate

IHC 1:20-1:200

**Basic Information** 

**Applications** 

Catalog Number: 14755-1-AP Size:

550 μg/ml Source: Rabbit

Isotype: IgG

Immunogen Catalog Number:

AG6503

**Tested Applications:** FC, IHC, IP, WB, ELISA Cited Applications:

IHC, WB

Species Specificity: human, mouse, rat **Cited Species:** human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

GenBank Accession Number: **Purification Method:** 

BC000408 GeneID (NCBI):

**UNIPROT ID:** Q9BWD1 Full Name:

acetyl-Coenzyme A acetyltransferase

Calculated MW: 41 kDa

Observed MW: 40-42 kDa

Positive Controls:

WB: Caco-2 cells, HepG2 cells, K-562 cells, MOLT-4

cells, mouse liver, rat liver IP: mouse liver tissue.

IHC: human liver tissue, human lung tissue, human

placenta tissue

## **Background Information**

## Notable Publications

Author	Pubmed ID	Journal	Application
De Huang	27644987	Cell Res	WB
George D Dalton	34568800	iScience	WB
Yunxia Yang	31679333	J Agric Food Chem	WB

Storage


Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


#### **Selected Validation Data**




WB result of ACAT2 antibody (14755-1-AP; 1:8000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ACAT2 transfected HepG2 cells.




Immunohistochemical analysis of paraffinembedded human liver using 14755-1-AP (ACAT2 antibody) at dilution of 1:100 (under 10x lens).




Immunohistochemical analysis of paraffinembedded human liver using 14755-1-AP (ACAT2 antibody) at dilution of 1:100 (under 40x lens).



IP result of anti-ACAT2 (IP:14755-1-AP, 4ug; Detection:14755-1-AP 1:1000) with mouse liver tissue lysate 4000ug.



1X10^6 HepG2 cells were stained with 0.2ug ACAT2 antibody (14755-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1500.



Various lysates were subjected to SDS PAGE followed by western blot with 14755-1-AP (ACAT2 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.