

For Research Use Only

AK1 Polyclonal antibody

Catalog Number: 14978-1-AP

4 Publications



Basic Information

Catalog Number:

14978-1-AP

Size:

400 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG6890

GenBank Accession Number:

BC001116

GeneID (NCBI):

203

UNIPROT ID:

P00568

Full Name:

adenylate kinase 1

Calculated MW:

22 kDa

Observed MW:

22 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

IHC 1:50-1:500

Applications

Tested Applications:

IHC, WB, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : mouse heart tissue, human heart tissue, mouse skeletal muscle tissue, rat heart tissue

IHC : mouse heart tissue,

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Gerald Wölkart	35618039	Eur J Pharmacol	WB
Michiyo Maruyama	34982774	PLoS One	WB
Yue Zong	30948787	Cell Res	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Immunohistochemical analysis of paraffin-embedded mouse heart tissue slide using 14978-1-AP (AK1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse heart tissue slide using 14978-1-AP (AK1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



mouse heart lysates were subjected to SDS PAGE followed by western blot with 14978-1-AP (AK1 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.