

For Research Use Only

SC65 Polyclonal antibody

Catalog Number: 15288-1-AP

Featured Product

7 Publications



Basic Information

Catalog Number:

15288-1-AP

Size:

600 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG7480

GenBank Accession Number:

BC001047

GeneID (NCBI):

10609

UNIPROT ID:

Q92791

Full Name:

synaptonemal complex protein SC65

Calculated MW:

50 kDa

Observed MW:

50 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:8000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

Applications

Tested Applications:

IP, WB, ELISA

Cited Applications:

IF, IHC, WB

Species Specificity:

human, mouse

Cited Species:

human, mouse

Positive Controls:

WB : HeLa cells,

IP : mouse bladder tissue,

Background Information

SC65, also named as Leprecan-like protein 4 or Nucleolar autoantigen No55, is a 437 amino acid protein, which belongs to the leprecan family. SC65 is a protein of the interphase nucleolus and tumour-associated autoantigen in patients with prostate cancer.

Notable Publications

Author	Pubmed ID	Journal	Application
Masahiko Terajima	31173582	PLoS Genet	WB
Chen Fang	35805016	Cancers (Basel)	WB
Sarah M Zimmerman	30417103	AIMS Genet	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



IP result of anti-SC65 (IP:15288-1-AP, 4ug; Detection:15288-1-AP 1:500) with mouse bladder tissue lysate 3200ug.



WB result of SC65 antibody (15288-1-AP; 1:4000; incubated at room temperature for 1.5 hours) with sh-Control and sh-SC65 transfected HeLa cells.



Various lysates were subjected to SDS PAGE followed by western blot with 15288-1-AP (SC65 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.