For Research Use Only

## CPSF6 Polyclonal antibody

Catalog Number:15489-1-AP

Featured Product

9 Publications




Basic Information	Catalog Number: 15489-1-AP	GenBank Accession Number: BC000714		Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI): 11052		Recommended Dilutions: WB 1:500-1:1000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:250-1:1000 IF 1:50-1:500	
	300 µg/ml				
	Source: UNIPROT ID: Rabbit 016630				
	Rabbit	Full Name: cleavage and polyadenylation specific factor 6, 68kDa Calculated MW: 59 kDa			
	Isotype: IgG Immunogen Catalog Number:				
	AG7852				
		Observed MV 55-68 kDa	Observed MW: 55-68 kDa		
Applications	Tested Applications:		Positive Controls:		
	IF/ICC, IHC, IP, WB, ELISA Cited Applications:			8 : HeLa cells, HepG2 cells, Jurkat cells, PC-3 cells, DA-MB-231 cells, SW480 cells	
	WB		IP : HeLa cells,		
	human	numan kidney tissu		n liver tissue, human heart tissue, huma	
	Cited Species:				
	numan				
	Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0	atively, antige	th		
Background Information	Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 The binding of Cleavage factor In assembly of the cleavage and po factors. CFIM is required for the fi heterologously expressed 25- an subcellular localization [PMID:15	atively, antige d with citrate n (CFIM), also kno lyadenylation ma irst step in pre-mR d 68-kDa subunits. 5169763]. In additi oplasm and contain	th n wwn as CPSF6, to the pr chinery and facilitate NA 3' -end processing . It involved in RNA bi on, it is a pre-mRNA p ns a C-terminal nuclea	re-mRNA is one of the earliest steps in th s the recruitment of other processing g and can be reconstituted in vitro from i nding, protein-protein interactions, and	
	Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 The binding of Cleavage factor Ir assembly of the cleavage and po factors. CFIM is required for the fi heterologously expressed 25- an- subcellular localization [PMID:15 between the nucleus and the cytor	atively, antige d with citrate n (CFIM), also kno lyadenylation ma irst step in pre-mR d 68-kDa subunits. 5169763]. In additi oplasm and contain	th n wwn as CPSF6, to the pr chinery and facilitate NA 3' -end processing . It involved in RNA bi on, it is a pre-mRNA p ns a C-terminal nuclea	re-mRNA is one of the earliest steps in th s the recruitment of other processing g and can be reconstituted in vitro from i nding, protein-protein interactions, and rocessing protein that dynamically shut	
	Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 The binding of Cleavage factor Ir assembly of the cleavage and po factors. CFIM is required for the fi heterologously expressed 25- an subcellular localization [PMID:15 between the nucleus and the cyto domain of the type bound by TNP	n (CFIM), also kno lyadenylation ma irst step in pre-mR d 68-kDa subunits. 5169763]. In additi pplasm and contai 903[PMID:1516976	th n wwn as CPSF6, to the pr chinery and facilitate NA 3' -end processing . It involved in RNA bi on, it is a pre-mRNA p ns a C-terminal nuclea 53,19864460].	re-mRNA is one of the earliest steps in th s the recruitment of other processing g and can be reconstituted in vitro from i nding, protein-protein interactions, and rocessing protein that dynamically shut ar-targeting arginine/serine-rich (RS-)	
Background Information	Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 The binding of Cleavage factor In assembly of the cleavage and po factors. CFIM is required for the fi heterologously expressed 25- an subcellular localization [PMID:15 between the nucleus and the cytor domain of the type bound by TNP	n (CFIM), also kno lyadenylation ma irst step in pre-mRi d 68-kDa subunits. 5169763]. In additi oplasm and contain 'O 3[PMID:1516976	th n wwn as CPSF6, to the pr chinery and facilitate NA 3' -end processing . It involved in RNA bi on, it is a pre-mRNA p ns a C-terminal nucles 63,19864460]. Journal	re-mRNA is one of the earliest steps in the s the recruitment of other processing g and can be reconstituted in vitro from in nding, protein-protein interactions, and rocessing protein that dynamically shut ar-targeting arginine/serine-rich (RS-) Application WB	


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




HeLa cells were subjected to SDS PAGE followed by western blot with 15489-1-AP (CPSF6 antibody) at dilution of 1:100 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human kidney using 15489-1-AP (CPSF6 antibody) at dilution of 1:50 (under 10x lens).




Immunohistochemical analysis of paraffinembedded human kidney using 15489-1-AP (CPSF6 antibody) at dilution of 1:50 (under 40x lens).




IP result of anti-CPSF6 (IP:15489-1-AP, 4ug; Detection:15489-1-AP 1:300) with HeLa cells lysate 3200ug.



Various lysates were subjected to SDS PAGE followed by western blot with 15489-1-AP (CPSF6 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 15489-1-AP (CPSF6 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 15489-1-AP (CPSF6 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed SW480 cells using CPSF6 antibody (15489-1-AP) at dilution of 1:200 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594phalloidin (red).