

For Research Use Only

BAG1L-specific Polyclonal antibody



Catalog Number: 16148-1-AP **1 Publications**

Basic Information

Catalog Number: 16148-1-AP	GenBank Accession Number: NM_004323	Purification Method: Antigen affinity purification
Size: 300 µg/ml	GeneID (NCBI): 573	Recommended Dilutions: IF 1:10-1:100
Source: Rabbit	UNIPROT ID: Q99933	
Isotype: IgG	Full Name: BCL2-associated athanogene	
	Calculated MW: 39 kDa	

Applications

Tested Applications: IF/ICC, ELISA	Positive Controls: IF : HepG2 cells,
Species Specificity: human	

Background Information

BAG1 has been identified that modulates gene transcription through poorly defined mechanisms. Four isoforms of the BAG1 protein (BAG1S, BAG1, BAG1M, and BAG1L) can be produced from a common mRNA by use of alternative translation initiation sites, including a non-canonical CTG codon in one instance. The longest, BAG1L (Mr ~50K), contains a nuclear localization signal (NLS) and resides in the nucleus, whereas BAG1M (Mr ~46K) has an incomplete NLS and distributes mainly in cytosol, unless dragged into the nucleus through interactions with other. Distribution of BAG1S(p33) is not clear yet. This antibody can recognize BAG1L and BAG1V.

Notable Publications

Author	Pubmed ID	Journal	Application
Nada Lallous	27765852	Mol. Cancer Ther	


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of HepG2 cells, using BAG1 antibody 16148-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).