For Research Use Only

BAG1S/1M/1L Polyclonal antibody

Catalog Number: 16150-1-AP

Featured Product

1 Publications



Basic Information

Catalog Number: 16150-1-AP

Size: 550 μg/ml

Source: Rabbit Isotype:

IgG

GenBank Accession Number:

NM_004323 GeneID (NCBI):

UNIPROT ID: Q99933 Full Name:

BCL2-associated athanogene

Calculated MW: 26 kDa Observed MW:

46-50 kDa, 41-46 kDa, 25-33 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:500-1:1000

IHC 1:20-1:200

Danisius Camerala

Applications

Tested Applications: IHC, WB, ELISA Cited Applications:

IHC, WB

Species Specificity: mouse, human Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls

WB: mouse brain tissue, Jurkat cells, mouse liver

tissue

IHC : human testis tissue, human kidney tissue, human lung tissue, human ovary tissue, human placenta

tissue, human spleen tissue

Background Information

BAG1 have been identified that modulate gene transcription through poorly defined mechanisms. Four isoforms of the BAG1 protein (BAG1S, BAG1, BAG1M and BAG1L) can be produced from a common mRNA by use of alternative translation initiation sites, including a non-canonical CTG codon in one instance. The longest, BAG1L (Mr~50K), contains a nuclear localization signal (NLS) and resides in the nucleus, whereas BAG1M (Mr~46K) has an incomplete NLS and distributes mainly in cytosol, unless dragged into the nucleus through interactions with other. Distribution of BAG1S(p33) is not clear yet. This antibody recongizes all the isoforms of BAG1, includes GAG1L, BAG1V, BAG1M and BAG1S.

Notable Publications

| Author | Pubmed ID | Journal | Application |
|--------------|-----------|-------------|-------------|
| Wenbai Huang | 26846101 | Tumour Biol | WB,IHC |

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



mouse brain tissue were subjected to SDS PAGE followed by western blot with 16150-1-AP (BAG1S/1M/1L antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human testis using 16150-1-AP (BAG1S/1M/1L antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human testis using 16150-1-AP (BAG1S/1M/1L antibody) at dilution of 1:50 (under 40x lens).