For Research Use Only

BAG1S/1M/1L Polyclonal antibody

Catalog Number: 16150-1-AP

Featured Product

1 Publications



Basic Information

Catalog Number: 16150-1-AP

Size: 550 μg/ml

Source: Rabbit Isotype:

IgG

GenBank Accession Number:

NM_004323 GeneID (NCBI):

LINUDDOT

UNIPROT ID: Q99933 Full Name:

BCL2-associated athanogene

Calculated MW: 26 kDa Observed MW:

46-50 kDa, 41-46 kDa, 25-33 kDa

Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:1000 IHC 1:20-1:200

Applications

Tested Applications: WB, IHC, ELISA

Cited Applications:

WB, IHC

Species Specificity: mouse, human Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls

 $\label{eq:WB:mouse} \textbf{WB:} \ \textbf{mouse} \ \textbf{brain} \ \textbf{tissue, Jurkat} \ \textbf{cells, mouse} \ \textbf{liver}$

tissue

IHC : human testis tissue, human kidney tissue, human lung tissue, human ovary tissue, human placenta

tissue, human spleen tissue

Background Information

BAG1 has been identified that modulates gene transcription through poorly defined mechanisms. Four isoforms of the BAG1 protein (BAG1S, BAG1, BAG1M, and BAG1L) can be produced from a common mRNA by use of alternative translation initiation sites, including a non-canonical CTG codon in one instance. The longest, BAG1L (Mr~50K), contains a nuclear localization signal (NLS) and resides in the nucleus, whereas BAG1M (Mr~46K) has an incomplete NLS and distributes mainly in cytosol, unless dragged into the nucleus through interactions with other. Distribution of BAG1S(p33) is not clear yet. This antibody recognizes three isoforms of BAG1, includes GAG1L, BAG1M, and BAG1S.

Notable Publications

Author	Pubmed ID	Journal	Application
Wenbai Huang	26846101	Tumour Biol	WB,IHC

Storage

Storage:

Store at -20 $^{\circ}\text{C}$. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



mouse brain tissue were subjected to SDS PAGE followed by western blot with 16150-1-AP (BAG1S/1M/1L antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human testis using 16150-1-AP (BAG1S/1M/1L antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human testis using 16150-1-AP (BAG1S/1M/1L antibody) at dilution of 1:50 (under 40x lens).