For Research Use Only

PSPC1 Polyclonal antibody

Catalog Number: 16714-1-AP

6 Publications



Basic Information

Catalog Number: GenBank Accession Number: 16714-1-AP BC014184

Size: GeneID (NCBI): 55269

Source: UNIPROT ID:

Rabbit Q8WXF1
Isotype: Full Name:

gG paraspeckle component 1

Immunogen Catalog Number:Calculated MW:AG10136523 aa, 59 kDaObserved MW:

66 kDa

Purification Method: Antigen affinity purification Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

protein lysate IF 1:50-1:500

WB 1:5000-1:50000

Applications

Tested Applications: IF/ICC, IP, WB, ELISA

Cited Applications:

IF, WB

Species Specificity: human, mouse, rat Cited Species: human, mouse Positive Controls:

WB: HEK-293 cells, HepG2 cells, PC-3 cells

IP: HEK-293 cells, IF: PC-3 cells,

Background Information

PSPC1 (paraspeckle component 1), a known paraspeckle biomarker, is a putative transcription factor that belongs to the Drosophila behavior/human splicing (DBHS) family. PSPC1 was recently identified as a contextual determinant of tumor progression in multiple cancer types involving oncogenic reprogramming to switch proapoptotic TGF- β to prometastatic TGF- β via hijacking of Smad2/3 targeting. Upregulated PSPC1 is correlated with advanced tumor stages and poor survival of patients with breast, lung, and liver cancers. Upregulated PSPC1 potentiates expression of mesenchymal markers, EMT transcription factors (EMT-TF), cancer stem-like cell transcription factors (CSC-TF), and c-Myc-related proliferation genes, thus promoting migration, invasion, spheroid formation, tumor formation, and metastasis.

Notable Publications

Author	Pubmed ID	Journal	Application
Jing Fan	30032211	Nucleic Acids Res	WB
Ralf Kleene	36674445	Int J Mol Sci	WB
Wei Zhang	37647746	Drug Resist Updat	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 16714-1-AP (PSPC1 antibody) at dilution of 1:12000 incubated at room temperature for 1.5 hours.



IP result of anti-PSPC1 (IP:16714-1-AP, 4ug; Detection:16714-1-AP 1:1000) with HEK-293 cells lysate 2800ug.



Immunofluorescent analysis of (10% Formaldehyde) fixed PC-3 cells using 16714-1-AP (PSPC1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).