For Research Use Only

ACTA1-Specific Polyclonal antibody

Catalog Number:17521-1-AP

10 Publications



Basic Information	Catalog Number:		ccession Number:	Purification Method:	
	17521-1-AP	NM_001100		Antigen affinity purification	
	Size: 260 µg/ml	Genel D (NC 58	.BI):	Recommended Dilutions: WB 1:500-1:2000	
	Source: UNIPROT ID: Rabbit P68133):	IHC 1:50-1:500	
	Isotype: Full Name:				
	IgG actin, alpha 1, skeletal muscle				
	Calculated MW: 42 kDa		MW:		
		Observed M	1W:		
		39-42 kDa			
Applications	Tested Applications:		Positive Co	ntrols:	
	FC, IHC, WB, ELISA	WB, ELISA WB : human skeletal muscle tissue, mouse sk		skeletal muscle tissue, mouse skeletal	
	Cited Applications:			ue, rat skeletal muscle tissue	
	IF, WB		IHC : human skeletal muscle tissue, human		
	Species Specificity:	nysteronyona tissue			
	human, mouse, rat				
	Cited Species: human, rat, mouse				
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	expressed in all eukaryotic in adult skeletal muscle, wh	cells. The ACTA1 gene nich interacts with a va nino acids difference a	e encodes skeletal musc ariety of proteins to proc among the actin family	cell motility and are ubiquitously le alpha-actin, the principal actin isofo Juce the force for muscle contraction. Th members. This antibody is specific to th	
	A. (1	Pubmed ID	Journal	Application	
Notable Publications	Author				
Notable Publications	Author Hongyi Zhou	34502418	Int J Mol Sci	WB	
Notable Publications		34502418 34786834	Int J Mol Sci J Cell Mol Med	WB	
Notable Publications	Hongyi Zhou				

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human skeletal muscle tissue slide using 17521-1-AP (ACTA1-Specific Antibody) at dilution of 1:500 (under 10x lens). human skeletal muscle tissue were subjected to SDS PAGE followed by western blot with 17521-1-AP (ACTA1-Specific antibody) at dilution of 1:1000



Immunohistochemical analysis of paraffin-embedded human skeletal muscle tissue slide using 17521-1-AP (ACTA1-Specific Antibody) at dilution of 1:500 (under 40x lens).



1X10^6 C2C12 cells were intracellularly stained with 0.4 ug Anti-Human ACTA1-Specific (17521-1-AP) and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).