For Research Use Only

JAK2 Polyclonal antibody

Catalog Number: 17670-1-AP

Featured Product

81 Publications



Purification Method:

WB 1:500-1:1000

protein lysate

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number: 17670-1-AP BC039695
Size: GeneID (NCBI): 3717
Source: UNIPROT ID: Rabbit 060674

Isotype:Full Name:IgGJanus kinase 2 (a protein tyrosine

Immunogen Catalog Number: kinase)

AG11264 Calculated MW: 1132 aa, 130 kDa

Observed MW: 150 kDa

Applications

Tested Applications:
IHC, IP, WB,ELISA
Cited Applications:
WB, IP, IF, IHC
Species Specificity:
human, mouse
Cited Species:

human, rat, mouse, zebrafish, pig

Positive Controls:

WB: Jurkat cells, NIH/3T3 cells

IP: Jurkat cells,

Background Information

Janus kinases (JAKs) are non-receptor tyrosine kinases that are activated by cytokine binding to cytokine receptors resulting in stimulation of the JAK-STAT signaling pathway. Aberrant activation of this signaling pathway is associated with numerous immune disorders and cancer. This antibody can detect full-length band of 130 kDa, and another bands 150 kDa after posttranslational modification.

Notable Publications

Author	Pubmed ID	Journal	Application
Zelong Gong	36289622	Biomedicines	WB
Yuanyuan Ren	36127737	Cancer Med	WB
Shaofang Chen	28944901	Mol Med Rep	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 17670-1-AP (JAK2 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 17670-1-AP (JAK2 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Jurkat cells were subjected to SDS PAGE followed by western blot with 17670-1-AP (JAK2 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



IP result of anti-JAK2 (IP:17670-1-AP, 4ug; Detection:17670-1-AP 1:300) with Jurkat cells lysate 3800ug.