For Research Use Only

## BMP9 Polyclonal antibody

Catalog Number:17769-1-AP




Basic Information	Catalog Number: 17769-1-AP	GenBank Accession Number: BC074921	Purification Method: Antigen affinity purification	
	Size: 450 µg/ml	GenelD (NCBI): 2658 UNIPROT ID: Q9UK05	Recommended Dilutions: WB 1:200-1:1000 IHC 1:50-1:500	
	Source: Rabbit			
	lsotype: IgG Immunogen Catalog Number: AG12177	Full Name: growth differentiation factor 2		
		Calculated MW: 429 aa, 47 kDa		
		Observed MW: 30 kDa, 46 kDa		
Applications	Tested Applications: IHC, WB,ELISA	Positive Controls:		
	Species Specificity: human	IHC : hum	WB : LO2 cells, SMMC-7721 cells IHC : human liver cancer tissue, human hepatocirrhosis tissue	
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
	<i>·</i> · · ·			
Background Informatio	<ul> <li>buffer pH 6.0</li> <li>BMP9 is a member of the bone more proteins is characterized by a poly containing seven conserved cystee differentiation in both embryonic adult liver and in differentiation or associated with hereditary hemore and has a calculated molecular m of about 13 kD. Once secreted, BM</li> </ul>	rphogenetic protein (BMP) family and basic proteolytic processing site whic ine residues. The members of this far and adult tissues. Studies in rodents s if cholinergic central nervous system i rhagic telangiectasia. Human BMP9 pi ass of about 47 kD. Posttranslational p	uggest that this protein plays a role in the neurons. Mutations in this gene are recursor protein contains 429 amino acids processing results in a mature BMP9 protein ons or as BMP9 proregion complexes in	


 For technical support and original validation data for this product please contact:

 T: 4006900926
 E: Proteintech-CN@ptglab.com
 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data





LO2 cells were subjected to SDS PAGE followed by western blot with 17769-1-AP (BMP9 Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 17769-1-AP (BMP9 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 17769-1-AP (BMP9 Antibody) at dilution of 1:200 (under 40x lens).