For Research Use Only

CD61 / Integrin Beta 3 Polyclonal antibody



Catalog Number:18309-1-AP

41 Publications


Basic Information	Catalog Number: GenBank Accession Number:			Purification Method:	
	18309-1-AP	BC127667	Antigen affinity	purification	
	Size:	GenelD (NCBI):	Recommended D		
	450 μg/ml 3690			WB 1:1000-1:4000 IP 0.5-4.0 ug for 1.0-3.0 mg of total	
	Source: Rabbit	UNIPROT ID: P05106	protein lucate		
	Isotype: Full Name: IgG integrin, beta 3 (platelet glycoprotein Illa, antigen CD61)				
	Calculated MW: 87 kDa				
	Observed MW: 100 kDa				
Applications	Tested Applications:		Positive Controls:		
	WB, IP, ELISA Cited Applications:	WB : Human peripheral blood platelets, mouse sple tissue, rat spleen tissue		atelets, mouse spleen	
	WB, IP, IF, FC, IHC, CoIP	IP : mouse spleen tissue,			
	Species Specificity: human, mouse, rat				
	Cited Species: human, rat, mouse, Hamster	r			
Background Information	Integrin beta-3, also named as CD61 and GPIIIa, is a receptor for cytotactin, fibronectin, laminin, matrix metalloproteinase-2, osteopontin, osteomodulin, prothrombin, thrombospondin, vitronectin and von Willebrand factor. Integrin beta-3 recognize the sequence R-G-D in a wide array of ligands and the sequence H-H-L-G-G-G-A-K-Q-A-G-D-V in fibrinogen gamma chain. Following activation integrin beta-3 brings about platelet/platelet interaction through binding of soluble fibrinogen which leads to rapid platelet aggregation which physically plugs ruptured endothelial surface. In case of HIV-1 infection, the interaction with extracellular viral Tat protein seems to enhance angiogenesis in Kaposi's sarcoma lesions. Human integrin beta-3 has a calculated molecular weight of 87 kDa. As a result of glycosylation, the apparent molecular mass of integrin beta-3 is approximately 90-110 kDa.				
Notable Publications	Author	Pubmed ID Jou	rnal	Application	
	Yujia Xu	32999280 Cell	. Death Dis	WB	
	Chao Cheng	36121292 App	l Immunohistochem Mol Morphol	IHC	
	Yueqi Li	30378153 J Ce	ll Biochem	WB	
Storage	Storage: Store at -20°C. Stable for on	e year after shipment.			

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data

hours.




Human peripheral blood platelets were subjected to SDS PAGE followed by western blot with 18309-1-AP (CD61 / Integrin Beta 3 antibody) at dilution of 1:2000 incubated at room temperature for 1.5



IP result of anti-CD61 / Integrin Beta 3 (IP:18309-1-AP, 4ug; Detection:18309-1-AP 1:300) with mouse spleen tissue lysate 5200 ug. 100 LDa→ ← 75 kDa→ 60 kDa→ 45 kDa→ 25 kDa→ 15 kDa→ 10 kDa→

180 kDa→ 140 kDa→

human peripheral blood platelets were subjected to SDS PAGE followed by western blot with 18309-1-AP (CD61 / Integrin Beta 3 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



IP result of anti-CD61 / Integrin Beta 3 (IP:18309-1-AP, 4ug; Detection:18309-1-AP 1:500) with mouse spleen tissue lysate 1920 ug.