For Research Use Only

Nestin Polyclonal antibody

Catalog Number:19483-1-AP 121 Publications



Basic Information

Catalog Number: 19483-1-AP Source: Rabbit Isotype:

GenBank Accession Number: NM_006617 GeneID (NCBI): 10763 **UNIPROT ID:** P48681 Full Name: nestin

Calculated MW: 177 kDa Observed MW:

170-180 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB: 1:500-1:1000 IF/ICC: 1:50-1:500

FC (Intra): 0.40 ug per 10^6 cells in a 100 µl suspension

Applications

Tested Applications: WB, IF/ICC, FC (Intra), ELISA Cited Applications:

WB, IHC, IF Species Specificity: human, mouse, rat **Cited Species:**

human, mouse, rat, monkey, zebrafish

Positive Controls:

WB: mouse brain tissue, rat brain tissue IF/ICC: SH-SY5Y cells, U-251 cells FC (Intra): SH-SY5Y cells,

Background Information

Nestin, also named as NES and Nbla00170, is a class VI intermediate filament protein expressed in stem cells of the central nervous system (CNS) but not in mature CNS cells. Its expression is used extensively as a marker for neural lineage cells derived from human embryonic stem (hES) and induced pluripotent stem (iPS) cells. Nestin is also $expressed\ in\ non-neural\ stem\ cell\ populations\ such\ as\ pancreatic\ islet\ progenitors\ and\ hematopoietic\ progenitors.$ Nestin can be detected as 220-240 kDa and 240-280 kDa. Studies point toward nestin heavy posttranslational modifications as possibly serving particular functions in podocytes and in a few other particular cell types(PMID: 17210924).

Notable Publications

Author	Pubmed ID	Journal	Application
Chenlong Li	31558707	Cell Death Dis	WB
Lili Ma	36132985	J Immunol Res	IF
Jiansheng Chen	31511040	J Exp Clin Cancer Res	WB


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Mouse brain lysates were subjected to SDS PAGE followed by western blot with 19483-1-AP (Nestin antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed SH-SY5Y cells using Nestin antibody (19483-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (-20°C Ethanol) fixed U-251 cells using Nestin antibody (19483-1-AP) at dilution of 1:400 and CoraLite@488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).



1x10^6 SH-SY5Y cells were intracellularly stained with 0.4 ug Nestin Polyclonal antibody (19483-1-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.4 ug rabbit IgG isotype control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.