For Research Use Only

ADRA1A-Specific Polyclonal antibody **proteintech**®

Catalog Number:19777-1-AP

Featured Product

5 Publications




Basic Information	Catalog Number: GenBank Accession Number: 19777-1-AP NM_000680			Purification Method: Antigen affinity purification	
	Size: 260 µg/ml	Genel D (NCBI): 148		Recommended Dilutions: WB 1:500-1:2000	
	Source: UNIPROT ID: Rabbit P35348				
	Isotype: Full Name: IgG adrenergic, alpha-1A-, receptor				
	Calculated MW: 51 kDa				
		Observed MW: 52 kDa			
Applications	Tested Applications: FC, WB, ELISA		Positive Controls:		
	VB: mouse heart tissue, HeLa cells, mouse brain tissue Cited Applications: WB, IF, IHC				
	Species Specificity: human, mouse, rat Cited Species: human, rat, mouse				
					Background Information
Notable Publications	Author	Pubmed ID Jou	urnal	Application	
	Miles Frost	32971843 Ph	armaceuticals (Base	l) IHC,IF	
	Jun-Jie Wang	26545627 Va	scul Pharmacol	WB	
	Miaomiao Zhu	35252820 iSc	tience	WB	
Storage	Storage: Store at -20°C. Stable for or Storage Buffer: PBS with 0.02% sodium azi Aliquoting is unnecessary f	de and 50% glycerol pH 7.3.			


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




mouse brain tissue lysates were subjected to SDS PAGE followed by western blot with 19777-1-AP (ADRA1A-Specific antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours without or with blocking peptide (1.4 ug peptide was used to block the binding of 1.0 ug antibody).



WB result of ADRA1A-Specific antibody (19777-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ADRA1A-Specific transfected HeLa cells.



mouse heart tissue were subjected to SDS PAGE followed by western blot with 19777-1-AP (ADRA1A-Specific antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



1X10^6 HepG2 cells were stained with 0.2ug ADRA1A-Specific antibody (19777-1-AP, red) and control antibody (blue). Fixed with 4% PFA blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1500.