For Research Use Only

KI67 Polyclonal antibody

Catalog Number: 19972-1-AP

40 Publications




Basic Information	Catalog Number: 19972-1-AP			Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):		Recommended Dilutions:	
	150 µg/ml 4288 IHC 1:200-1:800		IHC 1:200-1:800		
	Source: UNIPROTID: Rabbit P46013				
	Isotype: Full Name: IgG antigen identified by monoclonal antibody Ki-67				
		Calculated MW: 359 kDa			
Applications	Tested Applications: IHC,ELISA		Positive Controls:		
	IIIC, LUSA IHC : human breast cancer tissue, Cited Applications: WB, IF, IHC				
	Species Specificity: human				
	Cited Species: human, mouse, rat				
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	MKI67 gene encodes a nuclear protein, antigen KI-67, which was identified by monoclonal antibody Ki-67. Expression of this antigen occurs preferentially during late G1, S, G2 and M phases of the cell cycle, while in cells in G0 phase the antigen cannot be detected (PMID: 8227122). The antigen Ki-67 is a human nuclear protein the expression of which is strictly associated with cell proliferation and which is widely used in routine pathology as a "proliferation marker" to measure the growth fraction of cells in human tumors (PMID: 8227122).				
Notable Publications	Author	Pubmed ID	Journal	Application	
	Musin Kelel	32888953	J Invest Dermatol	IHC	
	Kai Tie		FASEB J	IHC	
	Yuan Zhang	26487539	Sci Rep	IHC	


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 19972-1-AP (KI67 Antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieved with Sodium Citrate buffer (pH 6.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 19972-1-AP (KI67 Antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieved with Sodium Citrate buffer (pH 6.0).