## For Research Use Only

# LDHC-Specific Polyclonal antibody

Catalog Number:19989-1-AP 4 Publications



**Basic Information** 

Catalog Number: 19989-1-AP Size: 400 μg/ml Source:

Rabbit P07864 Full Name: Isotype: lactate dehydrogenase C

Calculated MW: 36 kDa Observed MW:

30-36 kDa

NM\_002301

**UNIPROT ID:** 

3948

GeneID (NCBI):

GenBank Accession Number:

**Purification Method:** Antigen affinity purification

Recommended Dilutions:

WB 1:200-1:1000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:200-1:800 IF/ICC 1:10-1:100

**Applications** 

**Tested Applications:** WB, IHC, IF/ICC, IP, ELISA Cited Applications:

WB, IHC, IF Species Specificity: human, mouse, rat **Cited Species:** mouse, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: human testis tissue, mouse brain tissue, mouse lung tissue, mouse testis tissue, A549 cells, rat testis tissue

IP: mouse testis tissue,

IHC: mouse testis tissue, human testis tissue, human

heart tissue

IF/ICC: Hela cells.

## **Background Information**

LDHC, also named as CT32 and LDH-X, belongs to the LDH/MDH superfamily and LDH family. LDHC is a possible role in sperm motility. LDHC catalyzes the reaction: (S)-lactate + NAD+ = pyruvate + NADH. LDHC is a testis-specific isoform. Abnormal LDHC expression is associated with several forms of cancer. The antibody has no cross reaction with LDHA and LDHB.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Yuping Chen	34262451	Front Pharmacol	WB
Li Yang	31332157	Med Sci Monit	IHC
Rie Iida-Norita	36464740	Andrology	WB,IHC

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


### Selected Validation Data




human testis tissue were subjected to SDS PAGE followed by western blot with 19989-1-AP (LDHC-Specific antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 19989-1-AP (LDHC-Specific antibody) at dilution of 1:400 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 19989-1-AP (LDHC-Specific antibody) at dilution of 1:400 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of Hela cells, using LDHC antibody 19989-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



IP result of anti-LDHC-Specific (IP:19989-1-AP, 4ug; Detection:19989-1-AP 1:500) with mouse testis tissue lysate 2400ug.



Various lysates were subjected to SDS PAGE followed by western blot with 19989-1-AP (LDHC-Specific antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.