

# NCOR1 Polyclonal antibody

Catalog Number: 20018-1-AP

4 Publications

## Basic Information

## Catalog Number:

20018-1-AP

## Size:

500 µg/ml

## Source:

Rabbit

## Isotype:

IgG

## GenBank Accession Number:

NM\_006311

## GeneID (NCBI):

9611

## UNIPROT ID:

O75376

## Full Name:

nuclear receptor co-repressor 1

## Calculated MW:

270 kDa

## Observed MW:

120 kDa

## Purification Method:

Antigen affinity purification

## Recommended Dilutions:

WB 1:500-1:2000

IHC 1:20-1:200

## Applications

## Tested Applications:

IHC, WB, ELISA

## Cited Applications:

WB, ChIP

## Species Specificity:

human, mouse, rat

## Cited Species:

human, mouse

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Positive Controls:

WB : HepG2 cells, K-562 cells

IHC : human cervical cancer tissue, rat brain tissue

## Background Information

NCOR1, also named as KIAA1047 and TRAC1, belongs to the N-CoR nuclear receptor corepressors family. NCO1 mediates transcriptional repression by certain nuclear receptors. It is part of a complex which promotes histone deacetylation and the formation of repressive chromatin structures which may impede the access of basal transcription factors. The antibody recognizes the 103 kDa isoform of NCOR1.

## Notable Publications

Author	Pubmed ID	Journal	Application
Evangelia Lekka	36572670	Nat Commun	WB
Kenneth Allen Dyar	30096135	PLoS Biol	ChIP
Yan-Hang Wang	38250049	Theranostics	WB

## Storage

## Storage:

Store at -20°C. Stable for one year after shipment.

## Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



HepG2 cells were subjected to SDS PAGE followed by western blot with 20018-1-AP (NCOR1 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human cervical cancer using 20018-1-AP (NCOR1 antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human cervical cancer using 20018-1-AP (NCOR1 antibody) at dilution of 1:100 (under 40x lens).