### For Research Use Only

# AQP1 Polyclonal antibody

Catalog Number: 20333-1-AP

54 Publications



**Basic Information** 

Catalog Number: 20333-1-AP

 Size:
 GeneID (NCBI):

 500 μ g/ml
 358

Source: UNIPROT ID: Rabbit P29972
Isotype: Full Name:

IgG aquaporin 1 (Colton blood group)

Immunogen Catalog Number:Calculated MW:AG14093269 aa, 29 kDaObserved MW:

25-28 kDa, 35-50 kDa

GenBank Accession Number:

BC022486

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:5000-1:50000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:3000-1:12000 IF 1:50-1:500

**Applications** 

Tested Applications: WB,IP,IHC,IF-P,ELISA Cited Applications: WB,IP,IHC,IF,FC Species Specificity:

human, mouse, rat Cited Species:

Cat, human, Goat, Horse, rat, mouse, pig, canine, bovine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse skeletal muscle tissue, human heart tissue, rat skeletal muscle, mouse kidney, rat kidney

IP: mouse kidney tissue, mouse skeletal muscle tissue

IHC: mouse kidney tissue, human kidney tissue, human breast cancer tissue

IF: human breast cancer tissue, mouse kidney tissue

# **Background Information**

AQP1 is a member of aquaporins (AQPs) that are small membrane-spanning proteins facilitating water transport. AQP1 is expressed in most tissues in the mammalian body. Alterations of AQP1 expression have been linked to variety of diseases, including cancer. The predicted molecular weight of AQP1 is around 28 kDa, while highly glycosylated form can also be observed around 35-50 kDa. (PMID:20965731,16508653, 1530176).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Jianping Zhang	31572217	Front Physiol	WB,IHC
Haiyan Fu, Yuan	34622165	iScience	IF
Yi Song	36316968	Cell Prolif	IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 20333-1-AP (AQP1 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using AQP1 antibody (20333-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




IP result of anti-AQP1 (IP:20333-1-AP, 4ug; Detection:20333-1-AP 1:40000) with mouse kidney tissue lysate 2160 ug.




Various lysates were subjected to SDS PAGE followed by western blot with 20333-1-AP (AQP1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 20333-1-AP (AQP1 antibody) at dilution of 1:6000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 20333-1-AP (AQP1 antibody) at dilution of 1:6000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse kidney tissue using AQP1 antibody (20333-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).