For Research Use Only

AKT3 Polyclonal antibody

Catalog Number:21641-1-AP

13 Publications




Basic Information	Catalog Number: 21641-1-AP	GenBank Accession Number: BC 121154	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	760 µg/ml	10000	WB 1:500-1:1000	
	Source:	UNIPROT ID:	IHC 1:50-1:500	
	Rabbit	Q9Y243 Full Name: v-akt murine thymoma viral oncogene homolog 3 (protein kinase B, gamma)		
	Isotype:			
	IgG Immunogen Catalog Number:			
	AG16298			
		Calculated MW: 465 aa, 54 kDa		
		Observed MW:		
		60 kDa		
Applications	Tested Applications:	Positive Controls:		
	IHC, WB, ELISA	WD. mouse brain ussue, rat brain ussue		
	Cited Applications: IHC, WB	IHC : human tonsillitis tissue,		
	Species Specificity: human, mouse, rat			
	Cited Species: human			
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
Background Information	called the AKT kinase, and which and angiogenesis. AKT3 is a key	regulate many processes including m modulator of several tumors like mela	-protein kinases (AKT1, AKT2 and AKT3) letabolism, proliferation, cell survival, gro anoma, glioma and ovarian cancer. Active h highest levels present in advanced-stage	
Notable Publications	Author	Pubmed ID Journal	Application	
	Yang Yang	31515938 J Cell Mol Med	WB	
	Yong-Xia Wang	30186040 Cancer Cell Int	WB	
	Wei Sang	27329362 Cell Immunol	WB	
Storage	Storage: Store at -20°C. Stable for one yea Storage Buffer: PBS with 0.02% sodium azide an Aliquoting is unnecessary for -20	d 50% glycerol pH 7.3.		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 21641-1-AP (AKT3 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 21641-1-AP (AKT3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).