## For Research Use Only

## Ataxin 2 Polyclonal antibody

Catalog Number:21776-1-AP

Featured Product

31 Publications

GenBank Accession Number:



**Basic Information** 

Catalog Number:

21776-1-AP BC114546 Size: GeneID (NCBI): 350 μg/ml 6311

Source: UNIPROT ID:
Rabbit Q99700

Isotype: Full Name:
IgG ataxin 2

Immunogen Catalog Number:Calculated MW:AG164701313 aa, 140 kDa

Observed MW: 140-150 kDa Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:2000-1:16000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

**Applications** 

Tested Applications: IHC, IP, WB, ELISA Cited Applications:

CoIP, IF, IHC, IP, WB Species Specificity: human, mouse, rat Cited Species:

human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HeLa cells, HEK-293 cells, Jurkat cells, Neuro-2a

cells

IP: HEK-293 cells,

IHC: mouse brain tissue, human gliomas tissue

**Background Information** 

ATXN2 contains a repeat structure with 22 or 23 triplets coding for glutamine and the (CAG)8CAA(CAG)4CAA(CAG)8 sequence; expansion of this domain to a size ≥34 triplets with a pure CAG sequence primarily causes autosomal dominant SCA2 [PMID:18418684], while ATXN2 expansions with CAA interruptions were observed as the cause of Levo-dopa responsive Parkinson's disease [PMID:10668726]. ATXN2 expansions associated with ALS were reported to be interrupted by at least one CAA triplet [PMID:21537950]

**Notable Publications** 

Author	Pubmed ID	Journal	Application
Garam Kim	36288714	Cell Rep	WB,IF
Lauren A Ostrowski	30417124	Commun Biol	WB
Gaston Bonenfant	30944179	J Virol	WB,IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 21776-1-AP (Ataxin 2 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 21776-1-AP (Ataxin 2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using Ataxin 2 antibody (21776-1-AP) at dilution of 1:900 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 21776-1-AP (Ataxin 2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-Ataxin 2 (IP:21776-1-AP, 4ug; Detection:21776-1-AP 1:1000) with HEK-293 cells lysate 1200ug.