For Research Use Only

## CEP152 Polyclonal antibody

Catalog Number:21815-1-AP

Featured Product

3 Publications



| Basic Information    | Catalog Number:<br>21815-1-AP  | GenBank Acc<br>NM_014985  | ession Number:            | Purification Method:<br>Antigen affinity purification |  |
|----------------------|--|---|---------------------------|---|--|
|                      | Size:  | GenelD (NCB   | I):                       | Recommended Dilutions:                                |  |
|                      | 350 µg/ml  | 22995   |                           | WB 1:200-1:1000                                       |  |
|                      | Source:<br>Rabbit  | ource. UNIPROTID.   |                           | IHC 1:50-1:500  |  |
|                      | Isotype:Full Name:IgGcentrosomal protein 152kDa                        |   |                           |   |  |
|                      |  | Calculated M<br>189 kDa   | Calculated MW:<br>189 kDa |   |  |
|                      |  | Observed MW:<br>66 kDa  |                           |   |  |
| Applications         | Tested Applications:   |   | Positive Controls:        |   |  |
|                      | IHC, WB, ELISA   |   | WB: mouse brain tissue,   |   |  |
|                      | Cited Applications:<br>IF, WB  |   | IHC : human brain tissue, |   |  |
|                      | Species Specificity:<br>human, mouse<br>Cited Species:<br>human, mouse |   |                           |   |  |
|                      |  |   |                           |   | Note-IHC: suggested antigen retrieval with<br>TE buffer pH 9.0; (*) Alternatively, antigen<br>retrieval may be performed with citrate<br>buffer pH 6.0 |
|                      | Background Information   | CEP152, also named as KIAA0912, is a regulator of genomic integrity and cellular response to DNA damage actin<br>through ATR-mediated checkpoint signaling. It is necessary for centrosome duplication. CEP152 functions as a<br>molecular scaffold facilitating the interaction of PLK4 and CENPJ, two molecules involved in centriole formatior<br>CEP152, ATR and CENPJ control cellular responses to DNA damage. CEP152 has some isoforms with MW 189 kDa<br>152 kDa, 110 kDa and 66 kDa. |                           |   |  |
| Notable Publications | Author   | Pubmed ID   | Journal                   | Application   |  |
|                      | Zhenguang Ying   | 36280838  | BMC Biol                  | WB,IF   |  |
|                      |  | 36395215  | Science                   | IF  |  |
|                      | Tianyu Wu  | 50575225  |                           |   |  |
| Notable Fublications | Tianyu Wu<br>I-Ling Lu   | 29950674  | Nat Commun                | WB,IF   |  |


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



Immunohistochemical analysis of paraffinembedded human brain tissue slide using 21815-1-AP (CEP152 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



mouse brain tissue were subjected to SDS PAGE followed by western blot with 21815-1-AP (CEP152 Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



IP result of anti-CEP152 (IP:21815-1-AP, 4ug; Detection:21815-1-AP 1:2000) with mouse brain tissue lysate 1280 ug.



Immunohistochemical analysis of paraffinembedded human brain tissue slide using 21815-1-AP (CEP152 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).