

MST1 Polyclonal antibody

Catalog Number: 22245-1-AP

Featured Product

26 Publications

Basic Information

Catalog Number:

22245-1-AP

Size:

800 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG17738

GenBank Accession Number:

BC093768

GeneID (NCBI):

6789

UNIPROT ID:

Q13043

Full Name:

serine/threonine kinase 4

Calculated MW:

487 aa, 56 kDa

Observed MW:

56 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:6000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF 1:20-1:200

Applications

Tested Applications:

IF/ICC, IHC, IP, WB, ELISA

Cited Applications:

WB, IP, IHC, IF

Species Specificity:

human, mouse, rat

Cited Species:

human, rat, sheep, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, HepG2 cells, rat liver tissue, Jurkat cells, Ramos cells, C2C12 cells, mouse spleen tissue, rat spleen tissue

IP: HeLa cells,

IHC: human prostate cancer tissue,

IF: HepG2 cells,

Background Information

Mammalian STE20-like serine-threonine kinase MST1, encoded by the STK4 gene, is a multifunctional protein. MST1 and its closest paralogs MST2 (encoded by the STK3 gene), MST3, and MST4 are members of the Class II Germinal Center Family of Protein Kinases. STK3/4 and LATS1/2 (large tumor suppressor 1 and 2) are core kinase components of the Hippo tumor suppressor pathway in mammals. In the conventional Hippo pathway, the STK3/4 and LATS1/2 signaling cascade phosphorylates and inactivates the transcriptional coactivator YAP1 (yes associated protein 1) and its close paralog WWTR1. YAP1 and WWTR1 do not have DNA binding domains and they exert their biological outputs, such as cell proliferation and survival, by interacting with the TEAD1-4 transcription factors. Lines of evidence have indicated that dysregulation or loss of STK4/Hippo signaling is linked to developmental disorders and carcinogenesis with poor prognosis. STK4 is a stress-induced kinase and it can be activated in response to cell-death inducers. Autophosphorylation of STK4 at Thr183 (Thr180 in STK3) in the activation loop is a key activation mechanism for STK4/3 because phosphorylation of Thr183/180 causes the cleavage of STK4 by caspases under apoptotic conditions. The caspase-cleavage results in a more active STK4 protein (STK4-N, an amino-terminally truncated STK4), which localizes into the nucleus and induces apoptosis through histone modifications and chromatin condensations.

Notable Publications

Author	Pubmed ID	Journal	Application
Ting-Ting Zhang	28893403	Tissue Cell	WB, IHC
Qingxu Liu	34678143	Cell	IF, WB
Wenjiao Li	30315205	Cell Death Differ	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of MST1 antibody (22245-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-MST1 transfected HepG2 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 22245-1-AP (MST1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human prostate cancer tissue slide using 22245-1-AP (MST1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human prostate cancer tissue slide using 22245-1-AP (MST1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of HepG2 cells using 22245-1-AP (MST1 antibody) at dilution of 1:50 and and Rhodamine-labeled goat anti-rabbit IgG (red).



IP result of anti-MST1 (IP:22245-1-AP, 4ug; Detection:22245-1-AP 1:1000) with HeLa cells lysate 2000ug.