

For Research Use Only

VAMP7/TI-VAMP Polyclonal antibody



Catalog Number: 22268-1-AP

Featured Product

5 Publications

Basic Information

Catalog Number:

22268-1-AP

Size:

1000 μ g/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_005638

GeneID (NCBI):

6845

UNIPROT ID:

P51809

Full Name:

vesicle-associated membrane protein 7

Calculated MW:

25 kDa

Observed MW:

20-25 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:8000

IP 0.5-4.0 μ g for 1.0-3.0 mg of total protein lysate

IHC 1:20-1:200

IF 1:10-1:100

Applications

Tested Applications:

IF/ICC, IHC, IP, WB, ELISA

Cited Applications:

WB, IF, Blocking

Species Specificity:

human, mouse, rat

Cited Species:

human, rat, mouse

Positive Controls:

WB: mouse brain tissue, rat brain tissue

IP: mouse brain tissue,

IHC: human liver tissue, human small intestine tissue

IF: HepG2 cells, HeLa cells

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

VAMP7, also named as SYBL1 and TI-VAMP, belongs to the synaptobrevin family. It is involved in the targeting and/or fusion of transport vesicles to their target membrane during transport of proteins from the early endosome to the lysosome. VAMP7 is required for heterotypic fusion of late endosomes with lysosomes and homotypic lysosomal fusion. It is required for calcium regulated lysosomal exocytosis. VAMP7 is involved in the export of chylomicrons from the endoplasmic reticulum to the cis Golgi. It is required for exocytosis of mediators during eosinophil and neutrophil degranulation, and target cell killing by natural killer cells. It is also required for focal exocytosis of late endocytic vesicles during phagosome formation. The antibody is specific to VAMP7.

Notable Publications

Author	Pubmed ID	Journal	Application
Jia-min Yan	34747299	Autophagy	WB
Jia Liu	31062916	Small	WB
Dongdong Wang	32042550	Adv Sci (Weinh)	Blocking

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 22268-1-AP (VAMP7/TI-VAMP antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human liver using 22268-1-AP (VAMP7/TI-VAMP antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human liver using 22268-1-AP (VAMP7/TI-VAMP antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of HepG2 cells using 22268-1-AP (VAMP7/TI-VAMP antibody) at dilution of 1:25 and Rhodamine-Goat anti-Rabbit IgG.



IP result of anti-VAMP7/TI-VAMP (IP:22268-1-AP, 4ug; Detection:22268-1-AP 1:1000) with mouse brain tissue lysate 3440ug.