### For Research Use Only

# Angiopoietin 1 Polyclonal antibody

Catalog Number:23302-1-AP

Featured Product

27 Publications

GenBank Accession Number:



**Basic Information** 

Catalog Number: 23302-1-AP

23302-1-AP BC152411
Size: GeneID (NCBI):

500 μ g/ml 28

Source: UNIPROT ID:
Rabbit Q15389
Isotype: Full Name:
IgG angiopoietin 1
Immunogen Catalog Number: Calculated MW:

AG18829 498 aa, 58 kDa Observed MW:

70 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:500-1:2000

IF 1:50-1:500

**Applications** 

Tested Applications: IF/ICC, WB, ELISA Cited Applications:

WB, IF

Species Specificity: human, mouse, rat Cited Species: human, rat, mouse

#### Positive Controls:

WB: HUVEC cells, mouse liver tissue, HepG2 cells, mouse heart tissue, A375 cells, rat heart tissue, HeLa cells, K-562 cells, C6 cells, U-87 MG cells

IF: HeLa cells,

# **Background Information**

Angiopoietin 1 is a 70-kDa secreted glycoprotein generated from vascular mural cells, pericytes, and certain other cells. Angiopoietin 1 participates in vascular differentiation through angiogenesis, which is the process of the growth and remodelling of existing vessels. Angiopoietin 1 is also involved in the maintenance and turnover of blood vessels in mature animals. Angiopoietin 1 binds to and activates the TEK/TIE2 receptor by inducing its dimerization and tyrosine phosphorylation, playing an important role in the regulation of angiogenesis. Overexpression of Angiopoietin 1 has been proven to occur in malignant glioblastoma, neuroblastoma, non-small cell lung cancer, and other tumors.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Su-Wei Hu	36155178	Life Sci	WB
Daohai Qian	30359310	Stem Cell Res Ther	WB
Yu Wang	29061393	Neurosci Lett	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 23302-1-AP (Angiopoietin 1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed Hela cells using 23302-1-AP (Angiopoietin 1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).