For Research Use Only

MRP2 Polyclonal antibody

Catalog Number:24893-1-AP 23 Publications



Basic Information	Catalog Number: 24893-1-AP	GenBank Accession Number: BC 136419	Purification Meth Antigen affinity p		
	Size:	GenelD (NCBI):	Recommended Di		
	227 µg/ml	1244	WB 1:500-1:1000		
	Source: Rabbit	UNIPROT ID: Q92887	IHC 1:20-1:200		
	Isotype: IgG Immunogen Catalog Number: AG18694	Full Name: ATP-binding cassette, sub-family C (CFTR/MRP), member 2	ng cassette, sub-family C		
		Calculated MW: 1545 aa, 174 kDa			
		Observed MW: 190-250 kDa			
Applications	Tested Applications:	Positive Con	Positive Controls:		
	IHC, WB,ELISA	WB : HepG2 of	cells,		
	Cited Applications: WB, IHC	IHC : human	n small intestine tissue,		
	Species Specificity: human				
	Cited Species: human, rat, monkey				
	Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performe buffer pH 6.0	natively, antigen			
	Multi-drug resistance protein 2 (MRP2), also known as ABCC2, is an ATP binding cassette (ABC) transporter responsible for biliary excretion of xenobiotics, endobiotics, and their metabolites. Deficiency in ABCC2 results in the clinical disorder Dubin-Johnson syndrome. MRP2 is found to be expressed in a variety of human cancers, and i associated with resistance of tumor cells to various anticancer drugs including cisplatin. The predicted molecular weight of MRP2 is 174 kDa, while mature MRP2 usually has a slower migration around 190-250 kDa due to the glycosylation. (16434545, 18834541, 23045960)				
Background Information	associated with resistance of tur weight of MRP2 is 174 kDa, while	nor cells to various anticancer drugs inclue e mature MRP2 usually has a slower migra	• • •	redicted molecula	
	associated with resistance of tur weight of MRP2 is 174 kDa, while	nor cells to various anticancer drugs inclue e mature MRP2 usually has a slower migra	• • •	redicted molecula) kDa due to the	
	associated with resistance of tur weight of MRP2 is 174 kDa, while glycosylation. (16434545, 18834	nor cells to various anticancer drugs inclue e mature MRP2 usually has a slower migra 541, 23045960)	• • •	redicted molecula	
	associated with resistance of turr weight of MRP2 is 174 kDa, while glycosylation. (16434545, 18834 Author	nor cells to various anticancer drugs inclue e mature MRP2 usually has a slower migra 5541, 23045960) Pubmed ID Journal	ation around 190-250	redicted molecula) kDa due to the Application	
Background Information	associated with resistance of tur weight of MRP2 is 174 kDa, while glycosylation. (16434545, 18834 Author Zhenyu Wang	nor cells to various anticancer drugs inclue mature MRP2 usually has a slower migra (541, 23045960) Pubmed ID Journal 31588244 Theranostics	ation around 190-250	redicted molecula DkDa due to the Application IHC	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





HepG2 cells were subjected to SDS PAGE followed by western blot with 24893-1-AP (MRP2 antibody at dilution of 1:600 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human small intestine tissue slide using 24893-1-AP (MRP2 antibody at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human small intestine tissue slide using 24893-1-AP (MRP2 antibody at dilution of 1:50 (under 40x lens).