# For Research Use Only

# ATG2B Polyclonal antibody

Catalog Number: 25155-1-AP

Featured Product

16 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 25155-1-AP BC015016 GeneID (NCBI): Size: 500 ug/ml 55102 Source: **UNIPROT ID:** Rabbit Q96BY7 Full Name:

Antigen affinity purification Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500 IF/ICC 1:50-1:500

**Purification Method:** 

Isotype:

IgG ATG2 autophagy related 2 homolog B (S. cerevisiae) Immunogen Catalog Number:

AG18412 Calculated MW: 2078 aa, 233 kDa

Observed MW: 240-250 kDa

**Applications** 

**Tested Applications:** WB, IHC, IF/ICC, ELISA **Cited Applications:** WB, IF, CoIP Species Specificity:

human **Cited Species:** human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells,

IHC: human pancreas cancer tissue, human stomach

tissue

IF/ICC: HepG2 cells,

# **Background Information**

# Notable Publications

Author	Pubmed ID	Journal	Application
Jay Xiaojun Tan	36071159	Nature	WB
Jee-Eun Lee	34720018	Autophagy	WB
Ying Li	33124742	FASEBJ	WB

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

# **Selected Validation Data**




Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 25155-1-AP (ATG2B antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 25155-1-AP (ATG2B antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



HeLa cells were subjected to SDS PAGE followed by western blot with 25155-1-AP (ATG2B Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using ATG2B antibody (25155-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).