For Research Use Only

HYAL1 Polyclonal antibody

Catalog Number: 25179-1-AP



Basic Information

Catalog Number: 25179-1-AP Size:

600 µg/ml Source: Rabbit Isotype:

lgG hyaluronoglucosaminidase 1

Immunogen Catalog Number:Calculated MW:AG17943435 aa, 48 kDaObserved MW:

60 kDa

BC035695

GeneID (NCBI):

UNIPROT ID:

Q12794 Full Name:

GenBank Accession Number:

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:1000-1:8000

IHC 1:50-1:500

Applications

Tested Applications: IHC, WB, ELISA Species Specificity: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HCT 116 cells, HeLa cells, HepG2 cells, MCF-7

cells

IHC: mouse liver tissue,

Background Information

Hyaluronic acid (HA) is a glycosaminoglycan that is believed to have numerous important biologic functions, including modulation of cell proliferation, migration, and differentiation, as well as the regulation of extracellular water and protein homeostasis. It is also an integral structural component of cartilage and other tissues and acts as a lubricant in joints. Hyaluronidases are a family of enzymes that catalyse the degradation of HA. In humans, there are five functional hyaluronidases: HYAL1, HYAL2, HYAL3, HYAL4 and HYAL5 (also known as SPAM1 or PH-20); plus a pseudogene, HYAL6 (also known as HYALP1). HYAL-1 is present in many tissues and is predominantly found in the plasma and urine (PMID: 16600643). In addition to its function in normal cellular hyaluronan turnover, human HYAL1 is implicated in cancer proliferation, angiogenesis, and inflammatory diseases (PMID: 17503783).

Storage


Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 25179-1-AP (HYAL1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 25179-1-AP (HYAL1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 25179-1-AP (HYAL1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).