For Research Use Only

CD38 Polyclonal antibody

Catalog Number:25284-1-AP 6 Publications




Basic Information	Catalog Number: 25284-1-AP	GenBank Accession Numbe BC007964	r: Purification Method: Antigen affinity purification
	Size:	GeneID (NCBI):	Recommended Dilutions:
	800 µg/ml	952	WB 1:1000-1:6000
	Source: Rabbit	ENSEMBL Gene ID: ENSG00000004468	IHC 1:50-1:500
	lsotype: IgG	UNIPROT ID: P28907	
	Immunogen Catalog Number: AG17868	Full Name: CD38 molecule	
		Calculated MW: 300 aa, 34 kDa	
		Observed MW: 40-45 kDa	
Applications	Tested Applications:	Pos	itive Controls:
	FC, IHC, WB, ELISA	WB	Raji cells, HL-60 cells, Daudi cells, Ramos cells
	Cited Applications: WB, IF, IHC	IHC tiss	: human tonsillitis tissue, human appendicitis ue
	Species Specificity: human		
	Cited Species:		
	human		
	human Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0	atively, antigen	
Background Information	Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 CD38, also known as ADP-ribosyl cytoplasmic tail, a single membr glycosylation sites (PMID: 23191) catalyze synthesis of cyclic ADP r ribose to adenosine diphosphorit	atively, antigen d with citrate cyclase 1, is a type II transmem ane-spanning domain, and a C-t 35). The extracellular domain of ibose from nicotinamide adenir ose (PMID: 10636863). CD38 is e ed in diverse processes such as	expressed on a variety of hematopoietic and nor
	Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 CD38, also known as ADP-ribosyl cytoplasmic tail, a single membr glycosylation sites (PMID: 23191) catalyze synthesis of cyclic ADP r ribose to adenosine diphosphorit hematopoietic cells and is involv activation, and chemotaxis (PMID	atively, antigen d with citrate cyclase 1, is a type II transmem ane-spanning domain, and a C-t 55). The extracellular domain of ibose from nicotinamide adenir ose (PMID: 10636863). CD38 is e ed in diverse processes such as b: 25938500).	erminal extracellular region with four N- CD38 has bifunctional enzyme activities that le dinucleotide (NAD) and hydrolysis of cyclic Al expressed on a variety of hematopoietic and nor generation of calcium-mobilizing metabolites, o
	Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 CD38, also known as ADP-ribosyl cytoplasmic tail, a single membr glycosylation sites (PMID: 23192) catalyze synthesis of cyclic ADP r ribose to adenosine diphosphorit hematopoietic cells and is involv	atively, antigen d with citrate cyclase 1, is a type II transmem ane-spanning domain, and a C-t 55). The extracellular domain of ibose from nicotinamide adenir ose (PMID: 10636863). CD38 is o ed in diverse processes such as b: 25938500).	erminal extracellular region with four N- CD38 has bifunctional enzyme activities that le dinucleotide (NAD) and hydrolysis of cyclic AI expressed on a variety of hematopoietic and nor generation of calcium-mobilizing metabolites, or Application
Background Information	Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 CD38, also known as ADP-ribosyl cytoplasmic tail, a single membr glycosylation sites (PMID: 23191) catalyze synthesis of cyclic ADP r ribose to adenosine diphosphorit hematopoietic cells and is involv activation, and chemotaxis (PMID Author	atively, antigen d with citrate cyclase 1, is a type II transmem ane-spanning domain, and a C-t 55). The extracellular domain of ibose from nicotinamide adenir ose (PMID: 10636863). CD38 is e ed in diverse processes such as b: 25938500). Pubmed ID Journal	erminal extracellular region with four N- CD38 has bifunctional enzyme activities that le dinucleotide (NAD) and hydrolysis of cyclic AI expressed on a variety of hematopoietic and nor generation of calcium-mobilizing metabolites, or Application
	Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 CD38, also known as ADP-ribosyl cytoplasmic tail, a single membr glycosylation sites (PMID: 23191) catalyze synthesis of cyclic ADP r ribose to adenosine diphosphorit hematopoietic cells and is involv activation, and chemotaxis (PMID Author Hua Bai	atively, antigen d with citrate cyclase 1, is a type II transmem ane-spanning domain, and a C-t 35). The extracellular domain of ibose from nicotinamide adenir iose (PMID: 10636863). CD38 is e ed in diverse processes such as b: 25938500). Pubmed ID Journal 30326911 Cell Comm	erminal extracellular region with four N- CD38 has bifunctional enzyme activities that le dinucleotide (NAD) and hydrolysis of cyclic Al expressed on a variety of hematopoietic and nor generation of calcium-mobilizing metabolites, of hun Signal IF WB,IHC


 For technical support and original validation data for this product please contact:

 T: 4006900926
 E: Proteintech-CN@ptglab.com
 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data






Various lysates were subjected to SDS PAGE followed by western blot with 25284-1-AP (CD38 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 25284-1-AP (CD38 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 25284-1-AP (CD38 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10^6 HL-60 cells were stained with 0.2ug CD38 antibody (25284-1-AP, red) and control antibody (blue). Fixed with 4% PFA blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1500.