

For Research Use Only

Desmoplakin Polyclonal antibody

Catalog Number: 25318-1-AP

Featured Product

13 Publications



Basic Information

Catalog Number:

25318-1-AP

Size:

600 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG18065

GenBank Accession Number:

BC140802

GeneID (NCBI):

1832

UNIPROT ID:

P15924

Full Name:

desmoplakin

Calculated MW:

2871 aa, 332 kDa

Observed MW:

240-330 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:2000-1:16000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:2500-1:10000

IF 1:750-1:3000

Applications

Tested Applications:

FC, IF/ICC, IF-P, IHC, IP, WB, ELISA

Cited Applications:

IF, IP, WB

Species Specificity:

human, mouse, rat

Cited Species:

human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A431 cells, HepG2 cells, mouse heart tissue, BxPC-3 cells, HEK-293 cells

IP: BxPC-3 cells,

IHC: rat heart tissue, mouse heart tissue

IF: A431 cells, rat heart tissue

Background Information

The desmoplakin (DP) is the most abundant desmosomal proteins and is located at the cytoplasmic portion of the desmosomes which are specialized cell-cell adhesion structures abundant in tissues such as muscle and epidermis that are subjected to mechanical stress. Desmoplakin 1 and 2 (DP 1 and 2) are two splice variants sharing common C-terminal and N-terminal globular head and tail domains, but differ in the length of the rod domain that links them. This antibody detects both DP1 (280-330 kDa) and DP2 (240-260 kDa). (PMID: 16467215, 11781569)

Notable Publications

Author	Pubmed ID	Journal	Application
Jing Wang	36357564	Cancer Gene Ther	WB
Yanlei Yang	33993187	Cell Death Discov	WB
Tomohiko Sano	33802279	Cancers (Basel)	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of Desmoplakin antibody (25318-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-Desmoplakin transfected HepG2 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 25318-1-AP (Desmoplakin antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (-20°C Methanol) fixed A431 cells using Desmoplakin antibody (25318-1-AP) at dilution of 1:1500 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), Beta Actin antibody (66009-1-Ig, Clone: 2D4H5, red).




Immunohistochemical analysis of paraffin-embedded rat heart tissue slide using 25318-1-AP (Desmoplakin antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded rat heart tissue slide using 25318-1-AP (Desmoplakin antibody) at dilution of 1:5000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded mouse heart tissue slide using 25318-1-AP (Desmoplakin antibody) at dilution of 1:5000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).





Immunohistochemical analysis of paraffin-embedded mouse heart tissue slide using 25318-1-AP (Desmoplakin antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-Desmoplakin (IP:25318-1-AP, 4ug; Detection:25318-1-AP 1:10000) with BxPC-3 cells lysate 2080 ug.



Immunohistochemical analysis of paraffin-embedded mouse heart tissue slide using 25318-1-AP (Desmoplakin antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse heart tissue slide using 25318-1-AP (Desmoplakin antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

1X10⁶ A431 cells were intracellularly stained with 0.4 ug Anti-Human Desmoplakin (25318-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).