### For Research Use Only

# DGCR8 N-terminal Polyclonal antibody



Catalog Number: 25835-1-AP

**Featured Product** 

2 Publications

**Basic Information** 

Catalog Number: 25835-1-AP Size: 700 μg/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG22988

**Tested Applications:** 

Cited Applications:

Species Specificity:

**Cited Species:** bovine, mouse GenBank Accession Number:

BC078147 GeneID (NCBI): 54487 **UNIPROT ID:** Q8WYQ5 Full Name:

DiGeorge syndrome critical region

gene 8

Calculated MW: 773 aa, 86 kDa Observed MW: 88-100 kDa

**Purification Method:** 

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IF 1:20-1:200

**Applications** 

IF/ICC, IP, WB, ELISA

human

Positive Controls:

WB: HeLa cells, A431 cells

IP: A431 cells, IF: HeLa cells,

# **Background Information**

DGCR8 is a RNA-binding protein that assists the Rnase III enzyme Drosha in the processing of microRNAs (miRNAs), which regulate the expression of a large number of protein-coding genes[PMID: 22580560]. DGCR8, which contains two double-stranded RNA (dsRNA)-binding domains, may be an essential component of the primary miRNAs processing complex, along with Drosha, promoting the processing of primary microRNA to precursor microRNA. It is ubiquitous expressed in human and mouse tissues, and is deleted in DiGeorge syndrome[22323604]. The calculated molecular weight of DGCR8 is 82-86 kDa, but the post-modified DGCR8 is about 120 kDa.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
An Yan	36273819	Nucleic Acids Res	IF
Erika E Paulson	36322738	Proc Natl Acad Sci U S A	IF

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




WB result of DGCR8 antibody (25835-1-AP, 1:2000) with si-control and si-DGCR8 transfected A431 cells.



HeLa cells were subjected to SDS PAGE followed by western blot with 25835-1-AP (DGCR8 N-terminal antibody at dilution of 1:600 incubated at room temperature for 1.5 hours.



IP result of anti-DGCR8 N-terminal (IP:25835-1-AP, 4ug; Detection:25835-1-AP 1:600) with A431 cells lysate 2800ug.



Immunofluorescent analysis of (10% Formaldehyde) fixed HeLa cells using 25835-1-AP (DGCR8 N-terminal antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).