For Research Use Only

CEP83 Polyclonal antibody

Catalog Number: 26013-1-AP

8 Publications



Purification Method:

WB 1:500-1:1000 IHC 1:50-1:500

IF 1:200-1:800

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number:

26013-1-AP

BC125087

Size:

GeneID (NCBI):

600 µg/ml

51134

Source:

Rabbit

Q9Y592

Isotype:

GeneID (NCBI):

Full Name:

coiled-coil domain containing 41

Immunogen Catalog Number:Calculated MW:AG22826693 aa, 82 kDaObserved MW:

83 kDa, 68 kDa

Applications

Tested Applications: IF/ICC, IHC, WB, ELISA Cited Applications:

WB, IF

Species Specificity:

human
Cited Species:
human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HEK-293 cells,

IHC: human kidney tissue,

IF: hTERT-RPE1 cells,

Background Information

CEP83 (centrosomal protein, 83 kDa), also known as CCDC41 (coiled-coil domain containing 41), is required for ciliary vesicle docking to the mother centriole. It is a key component of the centriolar distal appendages. CCDC41 may collaborate with IFT20 in the trafficking of ciliary membrane proteins from the Golgi complex to the cilium during the initiation of primary cilium assembly (PMID: 23530209). Alternative splicing results in transcript variants encoding two isoforms with calculated molecular weights of 83 kDa and 68 kDa, respectively.

Notable Publications

Author	Pubmed ID	Journal	Application
Pierrick Le Borgne	36070319	PLoS Biol	IF
Min Liu	34480124	Cell Res	WB,IF
Delowar Hossain	31177295	Cell Mol Life Sci	IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




HEK-293 cells were subjected to SDS PAGE followed by western blot with 26013-1-AP (CEP83 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 26013-1-AP (CCDC41 antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 26013-1-AP (CCDC41 antibody) at dilution of 1:200 (under 40x lens).



Immunofluorescent analysis of (4% PFA) fixed hTERT-RPE1 cells using CEP83 antibody (26013-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CoraLite® 594 ARL13B antibody (CL594-17711, red).