### For Research Use Only

# CDK12/CRKRS Polyclonal antibody

Catalog Number: 26816-1-AP

Featured Product

**4 Publications** 



**Purification Method:** 

WB 1:2000-1:16000 IF 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

**Basic Information** 

**Applications** 

Catalog Number: 26816-1-AP Size: 1200 µg/ml

Rabbit Isotype:

Source:

Immunogen Catalog Number:

AG25323

Cited Applications:

Species Specificity:

**Tested Applications:** 

IF/ICC, WB, ELISA

human Cited Species: human

GenBank Accession Number:

BC150265 GeneID (NCBI): 51755 **UNIPROT ID:** Q9NYV4 Full Name:

Cdc2-related kinase, arginine/serine-

Calculated MW: 1490 aa, 164 kDa Observed MW: 180-200 kDa

Positive Controls:

WB: HeLa cells, K-562 cells, Jurkat cells

IF: HeLa cells,

## **Background Information**

CRKRS (Cdc2-related kinase, Arg/Ser), also named as cyclin-dependent kinase 12 (CKD12), belongs to CDC2/CDKX  $subfamily. \ CDK 12\ can form\ a\ heterodimeric\ complex\ with\ its\ activating\ partner,\ cyclin\ K,\ to\ regulate\ important$ cellular physiological processes that include transcriptional regulation, regulation of DNA damage repair genes such as BRCA1 and ATM, suppression of intronic polyadenylation, and susceptibility to PARP inhibitors (PMID: 30487607, 24240700, 31694772). CDK12 has 3 isoforms with the calculated molecular mass of 164/163/141 kDa, and the apparent molecular mass of 180-200 kDa due to some modifications (PMID 11683387, 22988298).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Yan Liang	34686673	Cell Death Discov	WB
Hui Li	33429148	Eur J Cancer	WB
Mareike Polenkowski	36590164	iScience	WB


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



WB result of CDK12/CRKRS antibody (26816-1-AP; 1:8000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CDK12/CRKRS transfected HeLa cells.



Various lysates were subjected to SDS PAGE followed by western blot with 26816-1-AP (CDK12/CRKRS antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (10% Formaldehyde) fixed HeLa cells using 26816-1-AP (CRKRS antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).