## For Research Use Only

# HNRNPM Polyclonal antibody

Catalog Number: 26897-1-AP

Featured Product

5 Publications



**Basic Information** 

Catalog Number: 26897-1-AP Size: 400 µg/ml Source: Rabbit

Immunogen Catalog Number:

AG25474

Isotype:

GenBank Accession Number:

BC000138 GeneID (NCBI): 4670 UNIPROT ID: P52272 Full Name:

heterogeneous nuclear ribonucleoprotein M Calculated MW:

78 kDa
Observed MW:
73-77 kDa

Purification Method:

Antigen affinity purification Recommended Dilutions: WB 1:1000-1:4000 IF 1:50-1:500

**Applications** 

Tested Applications: IF/ICC, WB, ELISA Cited Applications: IF, WB

Species Specificity: human

Cited Species: human, mouse

Positive Controls:

WB: COLO 320 cells, MCF-7 cells, HeLa cells, NIH/3T3

cells, Neuro-2a cells

IF: HepG2 cells,

# **Background Information**

HNRNPM, also named as NAGR1, is a 730 amino acid protein, which localizes in nucleus. HNRNPM as a pre-mRNA binding protein in vivo, binds avidly to poly(G) and poly(U) RNA homopolymers in vitro. HNRNPM is Involved in splicing. HNRNPM acts as a receptor for carcinoembryonic antigen in Kupffer cells, may initiate a series of signaling events leading to tyrosine phosphorylation of proteins and induction of IL-1 alpha, IL-6, IL-10 and tumor necrosis factor alpha cytokines. HNRNPM exists two isoforms with MV 77 kDa and 73 kDa.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Tao Wang	36151083	Nat Commun	IF
Wen-Juan Li	33782401	Nat Commun	WB
Pingfu Hou	31291975	J Exp Clin Cancer Res	WB

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


### **Selected Validation Data**




COLO 320 cells were subjected to SDS PAGE followed by western blot with 26897-1-AP (HNRNPM antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



WB result of HNRNPM antibody (26897-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-HNRNPM transfected MCF-7 cells.



COLO 320 cells were subjected to SDS PAGE followed by western blot with 26897-1-AP (HNRNPM antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using HNRNPM antibody (26897-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).