

For Research Use Only

# EMR1 Polyclonal antibody

Catalog Number: 27044-1-AP **29 Publications**



## Basic Information

<b>Catalog Number:</b> 27044-1-AP	<b>GenBank Accession Number:</b> BC059395	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 700 µg/ml	<b>GeneID (NCBI):</b> 2015	<b>Recommended Dilutions:</b> WB 1:500-1:2000 IHC 1:400-1:1600 IF 1:50-1:500
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> Q14246	
<b>Isotype:</b> IgG	<b>Full Name:</b> egf-like module containing, mucin-like, hormone receptor-like 1	
<b>Immunogen Catalog Number:</b> AG25883	<b>Calculated MW:</b> 886 aa, 97 kDa	
	<b>Observed MW:</b> 160 kDa	

## Applications

<b>Tested Applications:</b> IF/ICC, IHC, WB, ELISA	<b>Positive Controls:</b> WB : unboiled RAW 264.7 cells, HL-60 cells IHC : mouse spleen tissue, mouse liver tissue IF : mouse peritoneal macrophages,
<b>Cited Applications:</b> IF, IHC, WB	
<b>Species Specificity:</b> human, mouse	
<b>Cited Species:</b> human, rat, mouse, pig	
<b>Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b>	

## Background Information

EMR1 (EGF-like module containing mucin-like hormone receptor 1), also known as Adhesion G protein-coupled receptor E1 (ADGRE1), is a surface receptor with seven transmembrane segments that belong to the EGF-7-transmembrane family of G protein-coupled receptors (PMID: 14647991, 7601460). EMR1 expression is restricted to eosinophilic granulocytes, where expression is overlapping with the eotaxin receptor CCR3 and the immunoglobulin-like lectin Siglec-8. Absence on other leukocytes, including basophils, implies that EMR1 is a highly specific marker for eosinophils in humans and may be used as a novel therapeutic target for eosinophilic disorders (PMID: 17823986, 24530099). F4/80, the murine homolog of EMR1, is a marker of murine macrophage. The apparent molecular weight of F4/80 is 160 kDa, which is larger than the calculated molecular weight due to post-translational modifications (PMID: 7308288; 8647179).

## Notable Publications

Author	Pubmed ID	Journal	Application
Yu-Hui Gu	36386139	Front Pharmacol	IF
Junjie Li	33080309	Cancer Lett	IF
Huihui Tao	34843873	Toxicol Lett	IHC

## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




unboiled RAW 264.7 cells were subjected to SDS PAGE followed by western blot with 27044-1-AP (EMR1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse spleen tissue slide using 27044-1-AP (EMR1 antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed mouse peritoneal macrophages using EMR1 antibody (27044-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffin-embedded mouse spleen tissue slide using 27044-1-AP (EMR1 antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



HL-60 cells were subjected to SDS PAGE followed by western blot with 27044-1-AP (EMR1 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.