

For Research Use Only

Factor XII Polyclonal antibody

Catalog Number: 27154-1-AP



Basic Information

Catalog Number: 27154-1-AP	GenBank Accession Number: BC012390	Purification Method: Antigen affinity purification
Size: 800 µg/ml	GeneID (NCBI): 2161	Recommended Dilutions: WB 1:1000-1:3000 IHC 1:50-1:500
Source: Rabbit	UNIPROT ID: P00748	
Isotype: IgG	Full Name: coagulation factor XII (Hageman factor)	
Immunogen Catalog Number: AG25912	Calculated MW: 615 aa, 68 kDa	
	Observed MW: 80 kDa and 52 kDa	

Applications

Tested Applications:

IHC, WB, ELISA

Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: human plasma tissue, K-562 cells, HepG2 cells, Jurkat cells

IHC: human liver tissue,

Background Information

Factor XII (F XII, Hageman factor) is a 80kD, single chain glycoprotein that circulates in blood as an inactive zymogen. F XII plays an important role in blood coagulation, fibrinolysis, and kinin generation. Factor XII is activated by kallikrein in alpha-factor XIIa, which is further converted by trypsin into beta-factor XIIa. Alpha-factor XIIa is composed of a 52 kDa NH₂-terminal heavy chain, called coagulation factor XIIa heavy chain, and a 28 kDa COOH-terminal light chain, called coagulation factor XIIa light chain, connected by a disulfide bond. Beta-factor XIIa is composed of 2 chains linked by a disulfide bond, an N-terminal nonapeptide, called beta-factor XIIa part 1, and coagulation factor XIIa light chain, also known in this context as beta-factor XIIa part 2.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



human plasma were subjected to SDS PAGE followed by western blot with 27154-1-AP (F12 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 27154-1-AP (F12 antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 27154-1-AP (F12 antibody) at dilution of 1:200 (under 40x lens).