For Research Use Only

FBXW7 Polyclonal antibody

Catalog Number: 28424-1-AP

Featured Product

29 Publications



Basic Information

Catalog Number: GenBank Accession Number: 28424-1-AP NM_001349798 GeneID (NCBI): Size: 500 μg/ml 55294 **UNIPROT ID:** Source: Rabbit Q969H0 Full Name: Isotype:

F-box and WD repeat domain containing 7

Immunogen Catalog Number: AG28471 Calculated MW: 66 kDa

Observed MW:

100-110 kDa, 66-75 kDa

Applications

Tested Applications: IHC, WB, ELISA **Cited Applications:** WB, IP, IF, IHC Species Specificity: Human, Mouse **Cited Species:** human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HEK-293 cells, HepG2 cells, MCF-7 cells, mouse

Purification Method:

WB 1:500-1:2000 IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

IHC: human lung cancer tissue, human breast cancer

tissue

Background Information

FBXW7, also named as FBW7, FBX30, SEL10 and hAgo, is a substrate recognition component of a SCF (SKP1-CUL1-Fbox protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins. It probably recognizes and binds to phosphorylated target proteins. FBXW7 is involved in the degradation of cyclin-E, MYC, NOTCH1 released notch intracellular domain (NICD), and probably PSEN1. FBXW7 is a general tumor suppressor in human cancer (PMID: 17909001). FBXW7 has 3 isoforms (α / β / γ) $with the calculated \,molecular\,mass\,of\,80\,kDa,\,70\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa,\,and\,apparent\,molecular\,mass\,of\,100-110\,kDa\,and\,66\,kDa$ 66-75 kDa (PMID: 31346036/PMID: 22864569). K48 autoubiquitination is consistent with maintenance of ubiquitinated FBW7 (Ub-FBW7, 200 kDa)(PMID: 32353058).

Notable Publications

Author	Pubmed ID	Journal	Application
Chien-Hung Yeh	32907612	Mol Cancer	WB
Feng Qiu	34487730	Exp Cell Res	WB,IHC
Chongyang Li	33126914	J Exp Clin Cancer Res	WB,IHC

Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 28424-1-AP (FBXW7 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 28424-1-AP (FBXW7 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).