For Research Use Only

FANCD2 Polyclonal antibody

Catalog Number:28619-1-AP

Featured Product






Basic Information	Catalog Number: 28619-1-AP	GenBank Accession Number: NM_001018115	Purification Method: Antigen affinity purification	
	Size: 850 µg/ml	GeneID (NCBI): 2177	Recommended Dilutions: WB 1:500-1:2000	
	Source: Rabbit	UNIPROT ID: Q9BXW9		
	Isotype: IgG Immunogen Catalog Number: AG29460	Full Name: Fanconi anemia, complementation group D2 Calculated MW:		
	A029400	164 kDa		
		Observed MW: 150 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, ELISA Cited Applications: IF, IP, WB	WB : HeLa cells, K-562 cells		
	Species Specificity: Human			
	Cited Species: human			
Background Information	FANCD2, also named as FACD and FACD2, is required for maintenance of chromosomal stability. FANCD2 promote accurate and efficient pairing of homologs during meiosis. FANCD2 is involved in the repair of DNA double-strand breaks, both by homologous recombination and single-strand annealing. It may participate in S phase and G2 phase checkpoint activation upon DNA damage. It promotes BRCA2/FANCD1 loading onto damaged chromatin. FANCD2 may also be involved in B-cell immunoglobulin isotype switching. Defects in FANCD2 are a cause of Fanconi anemia (FA) which is a genetically heterogeneous, autosomal recessive disorder characterized by progressive pancytopenia, a diverse assortment of congenital malformations, and a predisposition to the development of malignancies.			
	Author	Pubmed ID Journal	Application	
Notable Publications	Zivian Huang	36382555 Adv Sci (Weinh)	WB	
Notable Publications	Zixian Huang			
Notable Publications		35754466 Front Pharmacol	WB,IP	
Notable Publications	Liying Song	35754466Front Pharmacol35489419Eur J Pharmacol	WB,IP IF	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 28619-1-AP (FANCD2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.