

For Research Use Only

Phospho-Beta Catenin (Ser33) Polyclonal antibody



Catalog Number: 28772-1-AP

1 Publications

Basic Information

Catalog Number:

28772-1-AP

Size:

400 µg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC058926

GeneID (NCBI):

1499

ENSEMBL Gene ID:

ENSG00000168036

UNIPROT ID:

P35222

Full Name:

catenin (cadherin-associated protein),
beta 1, 88kDa

Calculated MW:

781 aa, 86 kDa

Observed MW:

90-100 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB

Species Specificity:

Human

Cited Species:

human

Positive Controls:

WB : Calyculin A treated PC-3 cells, Calyculin A
treated HeLa cells

Background Information

Beta Catenin/ β -catenin, also known as CTNNB1, is a key downstream component of the canonical Wnt pathway that plays diverse and critical roles in embryonic development and adult tissue homeostasis. The N-terminal (1-49 amino acids) of β -catenin is the key region for its stability and the degradation by ubiquitin-proteasome system, and phosphorylation of β -catenin at the N-terminal Ser33, Ser37, Thr41, and Ser45 promotes its ubiquitination and degradation (PMID: 31930681). Phosphorylated β -catenin is targeted for ubiquitination-dependent proteolysis.

Notable Publications

Author	Pubmed ID	Journal	Application
Fei Zhou	37468964	Exp Hematol Oncol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated PC-3 cells and Calyculin A treated PC-3 cells were subjected to SDS PAGE followed by western blot with 28772-1-AP (Phospho-Beta Catenin (Ser33) antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH antibody as loading control.



Non-treated HeLa cells and Calyculin A treated HeLa cells were subjected to SDS PAGE followed by western blot with 28772-1-AP (Phospho-Beta Catenin (Ser33) antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH antibody as loading control.