For Research Use Only

CD45 Polyclonal antibody

Catalog Number: 29914-1-AP



Basic Information

Catalog Number:

29914-1-AP BC014239 Size: GeneID (NCBI): 500 μg/ml 5788

Source: ENSEMBL Gene ID:
Rabbit ENSG0000081237
Isotype: UNIPROT ID:
IgG P08575
Immunogen Catalog Number: Full Name:

AG31286 protein tyrosine phosphatase,

Calculated MW: 1304 aa, 147 kDa Observed MW: 180-250 kDa

receptor type, C

GenBank Accession Number:

Purification Method: Antigen affinity purification

Recommended Dilutions: WB 1:5000-1:50000 IHC 1:1000-1:4000

Applications

Tested Applications: IHC, WB, ELISA Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: Jurkat cells, Raji cells, Ramos cells

IHC: human tonsillitis tissue, human liver cancer

tissue, human colon tissue

Background Information

CD45, also known as leukocyte common antigen (LCA), B220, and T200, is a 180-240 kD single-chain type I membrane glycoprotein. It is a tyrosine phosphatase expressed on the plasma membrane of all hematopoietic cells, except erythrocytes and platelets. CD45 is a signaling molecule that regulates a variety of cellular processes including cell growth, differentiation, cell cycle, and oncogenic transformation. CD45 is one of the best negative selection markers for characterizing and/or isolating human mesenchymal stem cells from bone marrow and other sources. Along with the positive selection markers (such as CD29, CD44, CD90, CD105 and CD166), negative selection markers (such as CD34 and CD45) are used for MSC identification.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 29914-1-AP (CD45 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 29914-1-AP (CD45 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 29914-1-AP (CD45 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).